arXiv:1211.6395v3 [math.GR] 21 Dec 2012

Abstract

A countable group Γ is called shift-minimal if every non-trivial measure preserving action of Γ weakly contained in the Bernoulli shift $\Gamma \curvearrowright ([0,1]^\Gamma, \lambda^\Gamma)$ is free. We show that any group Γ whose reduced C^*-algebra $C^*_r(\Gamma)$ admits a unique tracial state is shift-minimal, and that any group Γ admitting a free measure preserving action of cost> 1 contains a finite normal subgroup N such that Γ/N is shift-minimal. Any shift-minimal group in turn is shown to have trivial amenable radical. Recurrence arguments are used in studying invariant random subgroups of a wide variety of shift-minimal groups. We also examine continuity properties of cost in the context of infinitely generated groups and equivalence relations. A number of open questions are discussed which concern cost, shift-minimality, C^*-simplicity, and uniqueness of tracial state on $C^*_r(\Gamma)$.

Shift-minimal groups, fixed price 1, and the unique trace property

Robin D. Tucker-Drob[‡]
Caltech[§]

December 21, 2012

Contents

1 Introduction **3**

2 Preliminaries **7**
2.1 Group theory 7
2.2 Ergodic theory 8
2.3 Invariant random subgroups 8

3 Shift-minimality **9**
3.1 Seven characterizations of shift-minimality 9
3.2 NA-ergodicity 10
3.3 Amenable invariant random subgroups 13

4 Permanence properties **14**
4.1 Invariant random subgroups with trivial intersection 14
4.2 Finite index subgroups 15
4.3 Direct sums 17
4.4 Other permanence properties 18

5 Examples of shift-minimal groups **19**
5.1 Free groups 19
5.2 Property (BP) 20
5.3 Linear groups 24
5.4 Unique tracial state on $C^*_r(\Gamma)$ 24

6 Cost **26**
6.1 Notation and background 26
6.2 Cost and weak containment in infinitely generated groups 27
6.3 The cost of a generic action 34

---

[*] **Keywords:** amenable radical, Bernoulli shift, C^*-simple, cost, fixed price, invariant random subgroup, non-free action, reduced C^*-algebra, stabilizer, strongly ergodic, unique trace, weak containment

[†] **MSC:** 37A15, 37A20, 37A25, 37A50, 37A55, 43A07, 60B99.

[‡] email:rtuckerd@caltech.edu

[§] Department of Mathematics; California Institute of Technology; Pasadena, CA 91125

6.4 Cost and invariant random subgroups . . . 35
6.5 Fixed price 1 and shift-minimality . . . 39

7 Questions **40**
7.1 General implications . . . 40
7.2 Cost and pseudocost . . . 41
7.3 Other questions . . . 42

Appendix A Invariant random subgroups as subequivalence relations **43**
A.1 Invariant random partitions . . . 43
A.2 Normalized subequivalence relations . . . 46

Appendix B The amenable radical of a countable group **49**
B.1 Basic properties of AR_Γ . . . 49
B.2 Groups with trivial amenable radical . . . 50

References **52**

1 Introduction

The *Bernoulli shift* of a countable discrete group Γ, denoted by s_Γ, is the measure preserving action $\Gamma \curvearrowright^{s} ([0,1]^\Gamma, \lambda^\Gamma)$ (where λ denotes Lebesgue measure on $[0,1]$) of Γ given by

$$(\gamma^s \cdot f)(\delta) = f(\gamma^{-1}\delta)$$

for $\gamma, \delta \in \Gamma$ and $f \in [0,1]^\Gamma$. If Γ is infinite, then the Bernoulli shift may be seen as the archetypal free measure preserving action of Γ. This point of view is supported by Abért and Weiss's result [AW11] that s_Γ is weakly contained in every free measure preserving action of Γ. Conversely, it is well known that any measure preserving action weakly containing a free action must itself be free. A measure preserving action is therefore free if and only if it exhibits approximate Bernoulli behavior.

Inverting our point of view, the approximation properties exhibited by s_Γ itself have been shown to reflect the group theoretic nature of Γ. One example of this is Schmidt's characterization [Sc81] of amenable groups as exactly those groups Γ for which s_Γ admits a non-trivial sequence of almost invariant sets. An equivalent formulation in the language of weak containment is that Γ is amenable if and only if s_Γ weakly contains an action that is not ergodic. In addition, a direct consequence of Foreman and Weiss's work [FW04] is that amenability of Γ is equivalent to *every* measure preserving action of Γ being weakly contained in s_Γ. That each of these properties of s_Γ is necessary for amenability of Γ is essentially a consequence of the Ornstein-Weiss Theorem [OW80], while sufficiency of these properties may be reduced to the corresponding representation theoretic characterizations of amenability due to Hulanicki and Reiter (see [Hu64, Hu66], [Zi84, 7.1.8], [BHV08, Appendix G.3]): a group Γ is amenable if and only if its left regular representation λ_Γ weakly contains the trivial representation if and only if λ_Γ weakly contains every unitary representation of Γ.

This paper investigates further the extent to which properties of a group may be detected by its Bernoulli action. Roughly speaking, it is observed that even when a group is non-amenable, the manifestation (or lack thereof) of certain behaviors in the Bernoulli shift has implications for the extent of that group's non-amenability. Central to this investigation is the following definition.

Definition 1.1. A countable group Γ is called *shift-minimal* if every non-trivial measure preserving action weakly contained in s_Γ is free.

The reader is referred to [Ke10] for background on weak containment of measure preserving actions. Note that by definition the trivial group $\{e\}$ is shift-minimal.

Shift-minimality, as with the above-mentioned ergodic theoretic characterizations of amenability, takes its precedent in the theory of unitary representations of Γ. It is well known that Γ is C^**-simple* (i.e., its reduced C^*-algebra $C^*_r(\Gamma)$ is simple) if and only if every non-zero unitary representation of Γ weakly contained in the left-regular representation λ_Γ is actually weakly equivalent to λ_Γ [Ha07]. Using the Abért-Weiss characterization of freeness it is apparent that Γ is shift-minimal if and only if every non-trivial m.p. action of Γ weakly contained in s_Γ is in fact weakly equivalent to s_Γ. Apart from analogy, the relationship between shift-minimality and C^*-simplicity in general is unclear. However, we show in Theorem 5.15 that shift-minimality follows from a property closely related to C^*-simplicity. A group Γ is said to have the *unique trace property* if there is a unique tracial state on $C^*_r(\Gamma)$.

Theorem 1.2. *Let Γ be a countable group. If Γ has the unique trace property then Γ is shift-minimal.*

In addition, a co-induction argument (Proposition 3.15) shows that shift-minimal groups have no non-trivial normal amenable subgroups, i.e., they have trivial amenable radical. This places shift-minimality squarely between two other properties whose general equivalence with C^*-simplicity remains an open problem. Indeed, it is open whether there are any general implications between C^*-simplicity and the unique trace property; in all concrete examples these two properties coincide. Furthermore, while the amenable radical of any C^*-simple group is known to be trivial [PS79], it is an open question – asked explicitly by Bekka and de la Harpe [BH00] – whether conversely, a group which is not C^*-simple always contains a non-trivial normal amenable subgroup. For shift-minimality in place of C^*-simplicity, a stochastic version of this question is shown to have a positive answer (Theorem 3.16).

Theorem 1.3. *A countable group Γ is shift-minimal if and only if there is no non-trivial amenable invariant random subgroup of Γ weakly contained in s_Γ.*

Here an *invariant random subgroup* (IRS) of Γ is a Borel probability measure on the compact space Sub_Γ of subgroups of Γ that is invariant under the conjugation action $\Gamma \curvearrowright \mathrm{Sub}_\Gamma$ of Γ. It is called *amenable* if it concentrates on the amenable subgroups of Γ. Invariant random subgroups generalize the notion of normal subgroups: if N is a normal subgroup of Γ then the Dirac measure δ_N on Sub_Γ is conjugation invariant. It is shown in [AGV12] that the invariant random subgroups of Γ are precisely those measures on Sub_Γ that arise as the stabilizer distribution of some measure preserving action of Γ (see §2.3).

Theorem 1.3 is not entirely satisfactory since it still seems possible that shift-minimality of Γ is equivalent to Γ having no non-trivial amenable invariant random subgroups whatsoever (see Question 7.4). In fact, the proof of Theorem 1.2 in §5.4 shows that this possibly stronger property is a consequence of the unique trace property.

Theorem 1.4. *Let Γ be a countable group. If Γ has the unique trace property then Γ has no non-trivial amenable invariant random subgroups.*

The known general implications among all of the notions introduced thus far are expressed in Figure 1.

Our starting point in studying shift-minimality is the observation that if $\Gamma \curvearrowright^a (X, \mu)$ is a m.p. action that is weakly contained in s_Γ then every non-amenable subgroup of Γ acts ergodically. We call this property of a m.p. action *NA-ergodicity.* We show in Theorem 3.13 that when a m.p. action of Γ is NA-ergodic then the stabilizer of almost every point must be amenable.

§4 deals with permanence properties of shift-minimality by examining situations in which freeness of a m.p. action $\Gamma \curvearrowright^a (X, \mu)$ may be deduced from freeness of some acting subgroup. Many of the proofs in this section appeal to some form of the Poincaré Recurrence Theorem.

A wide variety of groups are known to have the unique trace property and Theorem 1.2 shows that all such groups are shift-minimal. Among these are all non-abelian free groups ([Pow75]), all Powers groups



Figure 1: The solid lines indicate known implications and the dotted lines indicate open questions discussed in §7. Any implication which is not addressed by the diagram is open in general. However, these properties all coincide for large classes of groups, e.g., linear groups (see §5.3).

and weak Powers groups ([Ha85], [BN88]), groups with property P_{nai} [BCH94], all ICC relatively hyperbolic groups ([AM07]), and all ICC groups with a minimal non-elementary convergence group action [MOY11]. In §5 we observe that all of these groups share a common paradoxicality property we call (BP), abstracted from M. Brin and G. Picioroaga's proof that all weak Powers groups contain a free group (see [Ha07, following Question 15]). It is shown in Theorem 5.6 that any non-trivial ergodic invariant random subgroup of a group with property (BP) must contain a non-abelian free group almost surely. Recurrence once again plays a key role in the proof.

§6 studies the relationship between cost, weak containment, and invariant random subgroups. Kechris shows in [Ke10, Corollary 10.14] that if $\boldsymbol{a}$ and $\boldsymbol{b}$ are free measure preserving actions of a finitely generated group Γ then $\boldsymbol{a} \prec \boldsymbol{b}$ implies $C(\boldsymbol{b}) \leqslant C(\boldsymbol{a})$ where $C(\boldsymbol{a})$ denotes the cost of $\boldsymbol{a}$ (i.e., the cost of the orbit equivalence relation generated by $\boldsymbol{a}$). This is deduced from the stronger fact [Ke10, Theorem 10.13] that the cost function $C : \mathrm{FR}(\Gamma, X, \mu) \to \mathbb{R}$, $\boldsymbol{a} \mapsto C(\boldsymbol{a})$, is upper semi-continuous for finitely generated Γ. In §6.2 we obtain a generalization which holds for arbitrary countable groups (Theorem 6.4 below). The consequences of this generalization are most naturally stated in terms of an invariant we call *pseudocost*.

If E is a m.p. countable Borel equivalence relation on (X, μ) then the pseudocost of E is defined as $PC_\mu(E) := \mathbf{inf}_{(E_n)} \liminf_n C_\mu(E_n)$, where $(E_n)_{n\in\mathbb{N}}$ ranges over all increasing sequences $E_0 \subseteq E_1 \subseteq \cdots$, of Borel subequivalence relations of E such that $\bigcup_n E_n = E$. The pseudocost of an action and of a group is then defined in analogy with cost (see Definition 6.6). It is immediate that $PC_\mu(E) \leqslant C_\mu(E)$, and while the pseudocost and cost coincide in most cases, including whenever E is treeable or whenever $C_\mu(E) < \infty$ (Corollary 6.8), it is unclear whether equality holds in general.

One of the main motivations for introducing pseudocost is the following useful continuity property (Corollary 6.20):

Theorem 1.5. *Let $\boldsymbol{a} = \Gamma \curvearrowright^{a} (X, \mu)$ and $\boldsymbol{b} = \Gamma \curvearrowright^{b} (Y, \nu)$ be measure preserving actions of a countable group Γ. Assume that $\boldsymbol{a}$ is free. If $\boldsymbol{a} \prec \boldsymbol{b}$ then $PC(\boldsymbol{b}) \leqslant PC(\boldsymbol{a})$.*

Combining Theorem 1.5 and [AW11, Theorem 1] it follows that, among all free m.p. actions of Γ, the Bernoulli shift $\boldsymbol{s}_\Gamma$ has the maximum pseudocost. Since pseudocost and cost coincide for m.p. actions of finitely generated groups, this generalizes the result of [AW11] that $\boldsymbol{s}_\Gamma$ has the greatest cost among free actions of a finitely generated group Γ. In Corollary 6.22 we use Theorem 1.5 to deduce general consequences for cost:

Theorem 1.6. *Let $\boldsymbol{a}$ and $\boldsymbol{b}$ be m.p. actions of a countably infinite group Γ. Assume that $\boldsymbol{a}$ is free and $\boldsymbol{a} \prec \boldsymbol{b}$.*

1. *If* $\mathrm{C}(\boldsymbol{b}) < \infty$ *then* $\mathrm{C}(\boldsymbol{b}) \leqslant \mathrm{C}(\boldsymbol{a})$.
2. *If* E_b *is treeable then* $\mathrm{C}(\boldsymbol{b}) \leqslant \mathrm{C}(\boldsymbol{a})$.
3. *If* $\mathrm{C}(\boldsymbol{a}) = 1$ *then* $\mathrm{C}(\boldsymbol{b}) = 1$.

This leads to a characterization of countable groups with fixed price 1 as exactly those groups whose Bernoulli shift has cost 1. This characterization was previously shown for finitely generated groups in [AW11].

Theorem 1.7. *Let* Γ *be a countable group. The following are equivalent:*

(1) Γ *has fixed price* 1

(2) $\mathrm{C}(\boldsymbol{s}_\Gamma) = 1$

(3) $\mathrm{C}(\boldsymbol{a}) = 1$ *for some m.p. action* $\boldsymbol{a}$ *weakly equivalent to* $\boldsymbol{s}_\Gamma$.

(4) $\mathrm{PC}(\boldsymbol{a}) = 1$ *for some m.p. action* $\boldsymbol{a}$ *weakly equivalent to* $\boldsymbol{s}_\Gamma$.

(5) Γ *is infinite and* $\mathrm{C}(\boldsymbol{a}) \leqslant 1$ *for some non-trivial m.p. action* $\boldsymbol{a}$ *weakly contained in* $\boldsymbol{s}_\Gamma$.

We use this characterization to obtain a new class of shift-minimal groups in §6.5. In what follows, AR_Γ denotes the amenable radical of Γ (see Appendix B). Gaboriau [Ga00, Theorem 3] showed that if Γ does not have fixed price 1 then AR_Γ is finite. We now have:

Theorem 1.8. *Let* Γ *be a countable group that does not have fixed price* 1. *Then* AR_Γ *is finite and* $\Gamma/\mathrm{AR}_\Gamma$ *is shift-minimal.*

In Theorem 6.31 of §6.4 it is shown that if the hypothesis of Theorem 1.8 is strengthened to $\mathrm{C}(\Gamma) > 1$, i.e., if *all* free m.p. actions of Γ have cost > 1, then the conclusion may be strengthened considerably. The following is an analogue of Bergeron and Gaboriau's result [BG05, §5] (see also [ST10, Corollary 1.6]) in which the statement is shown to hold for the first ℓ^2-Betti number in place of cost.

Theorem 1.9. *Suppose that* $\mathrm{C}(\Gamma) > 1$. *Let* $\Gamma \curvearrowright^{a} (X, \mu)$ *be an ergodic measure-preserving action of* Γ *on a non-atomic probability space. Then exactly one of the following holds:*

1. *Almost all stabilizers are finite;*
2. *Almost every stabilizer has infinite cost, i.e.,* $\mathrm{C}(\Gamma_x) = \infty$ *almost surely.*

In particular, AR_Γ *is finite and* $\Gamma/\mathrm{AR}_\Gamma$ *has no non-trivial amenable invariant random subgroups.*

The analysis of pseudocost in §6.2 is used in §6.3 to study the cost of generic actions in the Polish space $A(\Gamma, X, \mu)$ of measure preserving actions of Γ. For any group Γ there is a comeager subset of $A(\Gamma, X, \mu)$, consisting of free actions, on which the cost function $\mathrm{C} : A(\Gamma, X, \mu) \to \mathbb{R} \cup \{\infty\}$ takes a constant value $\mathrm{C}_{\mathrm{gen}}(\Gamma) \in \mathbb{R}$ [Ke10]. Likewise, the pseudocost function $\boldsymbol{a} \mapsto \mathrm{PC}(\boldsymbol{a})$ must be constant on a comeager set of free actions, and we denote this constant value by $\mathrm{PC}_{\mathrm{gen}}(\Gamma)$. Kechris shows in [Ke10] that $\mathrm{C}_{\mathrm{gen}}(\Gamma) = \mathrm{C}(\Gamma)$ for finitely generated Γ and Problem 10.11 of [Ke10] asks whether $\mathrm{C}_{\mathrm{gen}}(\Gamma) = \mathrm{C}(\Gamma)$ in general. The following is proved in Corollaries 6.28 and 6.27.

Theorem 1.10. *Let* Γ *be a countably infinite group. Then*

1. *The set* $\{\boldsymbol{a} \in A(\Gamma, X, \mu) : \boldsymbol{a}$ *is free and* $\mathrm{PC}(\boldsymbol{a}) = \mathrm{PC}(\Gamma)\}$ *is dense* G_δ *in* $A(\Gamma, X, \mu)$.
2. $\mathrm{PC}_{gen}(\Gamma) = \mathrm{PC}(\Gamma)$.

3. *Either* $C_{gen}(\Gamma) = C(\Gamma)$ *or* $C_{gen}(\Gamma) = \infty$.
4. *If* $PC(\Gamma) = 1$ *then* $C_{gen}(\Gamma) = C(\Gamma) = 1$.
5. *The set*

$$\big\{\boldsymbol{b} \in A(\Gamma, X, \mu) : \boldsymbol{b} \textit{ is free and } \exists \textit{aperiodic Borel subequivalence relations} \\ E_0 \subseteq E_1 \subseteq E_2 \subseteq \cdots \textit{ of } E_b, \textit{ with } E_b = \bigcup_n E_n \textit{ and } \lim_n C_\mu(E_n) = C(\Gamma)\big\}$$

is dense G_δ *in* $A(\Gamma, X, \mu)$.

6. *If all free actions of* Γ *have finite cost then* $\{\boldsymbol{b} \in A(\Gamma, X, \mu) : \boldsymbol{b}$ *is free and* $C(\boldsymbol{b}) = C(\Gamma)\}$ *is dense* G_δ *in* $A(\Gamma, X, \mu)$.

The only possible exception to the equality $C_{\mathrm{gen}}(\Gamma) = C(\Gamma)$ would be a group Γ with $C(\Gamma) < \infty$ such that the set $\{\boldsymbol{a} \in A(\Gamma, X, \mu) : \boldsymbol{a}$ is free, $C(\boldsymbol{a}) = \infty$ and E_a is not treeable$\}$ comeager in $A(\Gamma, X, \mu)$.

A number of questions are discussed in §7. The paper ends with two appendices, the first clarifying the relationship between invariant random subgroups and subequivalence relations. The second contains relevant results about the amenable radical of a countable group.

Acknowledgments. I would like to thank Miklos Abért, Lewis Bowen, Clinton Conley, Yair Glasner, Alexander Kechris, and Jesse Peterson for valuable comments and suggestions. The research of the author was partially supported by NSF Grant DMS-0968710.

2 Preliminaries

Throughout, Γ denotes a countable discrete group. The identity element of Γ is denoted by e_Γ, or simply e when Γ is clear from the context. All countable groups are assumed to be equipped with the discrete topology; a *countable group* always refers to a countable discrete group.

2.1 Group theory

Subgroups. Let Δ and Γ be countable groups. We write $\Delta \leqslant \Gamma$ to denote that Δ is a subgroup of Γ and we write $\Delta \lhd \Gamma$ to denote that Δ is a normal subgroup of Γ. The index of a subgroup $\Delta \leqslant \Gamma$ is denoted by $[\Gamma : \Delta]$. The *trivial* subgroup of Γ is the subgroup $\{e_\Gamma\}$ that contains only the identity element. For a subset $S \subseteq \Gamma$ we let $\langle S \rangle$ denote the subgroup generated by S. A group that is not finitely generated will be called *infinitely generated.*

Centralizers and normalizers. Let S be any subset of Γ and let H be a subgroup of Γ. The *centralizer* of S in H is the set

$$C_H(S) = \{h \in H : \forall s \in S\ (hsh^{-1} = s)\}$$

and the *normalizer* of S in H is the set

$$N_H(S) = \{h \in H : hSh^{-1} = S\}.$$

Then $C_H(S)$ and $N_H(S)$ are subgroups of H with $C_H(S) \lhd N_H(S)$. Clearly $C_H(S) = C_\Gamma(S) \cap H$ and $N_H(S) = N_\Gamma(S) \cap H$. The group $C_\Gamma(\Gamma)$ is called the *center* of Γ and is denoted by $Z(\Gamma)$. We say that a subset T of Γ *normalizes* S if $T \subseteq N_\Gamma(S)$. We call a subgroup H *self-normalizing* in Γ if $H = N_\Gamma(H)$.

Infinite conjugacy class (ICC) groups. A group Γ is called *ICC* if every $\gamma \in \Gamma \setminus \{e\}$ has an infinite conjugacy class. This is equivalent to $C_\Gamma(\gamma)$ having infinite index in Γ for all $\gamma \neq e$. Thus, according to our definition, the trivial group $\{e\}$ is ICC.

The Amenable Radical. We let AR_Γ denote the *amenable radical* of Γ. See Appendix B below.

2.2 Ergodic theory

Measure preserving actions. A *measure preserving (m.p.) action* of Γ is a triple $(\Gamma, a, (X, \mu))$, which we write as $\Gamma \curvearrowright^a (X, \mu)$, where (X, μ) is a standard probability space (possibly with atoms) and $a : \Gamma \times X \to X$ is a Borel action of Γ on X that preserves the probability measure μ. For $(\gamma, x) \in \Gamma \times X$ we let $\gamma^a x$ denote the image $a(\gamma, x)$ of the pair (γ, x) under a. We write $\boldsymbol{a}$ for $\Gamma \curvearrowright^a (X, \mu)$ when Γ and (X, μ) are clear from the context. A measure preserving action of Γ will also be called a Γ-*system* or simply a *system* when Γ is understood.

For the rest of this subsection let $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ and let $\boldsymbol{b} = \Gamma \curvearrowright^b (Y, \nu)$.

Isomorphism and factors. If $\varphi : (X, \mu) \to Y$ is a measurable map then we let $\varphi_* \mu$ denote the pushforward measure on Y given by $\varphi_* \mu(B) = \mu(\varphi^{-1}(B))$ for $B \subseteq Y$ Borel. We say that $\boldsymbol{b}$ is a *factor* of $\boldsymbol{a}$ (or that $\boldsymbol{a}$ *factors onto* $\boldsymbol{b}$), written $\boldsymbol{b} \sqsubseteq \boldsymbol{a}$, if there exists a measurable map $\pi : X \to Y$ with $\pi_* \mu = \nu$ and such that for each $\gamma \in \Gamma$ the equality $\pi(\gamma^a x) = \gamma^b \pi(x)$ holds for μ-almost every $x \in X$. Such a map π is called a *factor map* from $\boldsymbol{a}$ to $\boldsymbol{b}$. The factor map π is called an *isomorphism* from $\boldsymbol{a}$ to $\boldsymbol{b}$ if there exists a co-null subset of X on which π is injective. We say that $\boldsymbol{a}$ and $\boldsymbol{b}$ are *isomorphic*, written $\boldsymbol{a} \cong \boldsymbol{b}$, if there exists some isomorphism from $\boldsymbol{a}$ to $\boldsymbol{b}$.

Weak containment of m.p. actions. We write $\boldsymbol{a} \prec \boldsymbol{b}$ to denote that $\boldsymbol{a}$ is weakly contained in $\boldsymbol{b}$ and we write $\boldsymbol{a} \sim \boldsymbol{b}$ to denote that $\boldsymbol{a}$ and $\boldsymbol{b}$ are weakly equivalent. The reader is referred to [Ke10] for background on weak containment of measure preserving actions.

Product of actions. The *product* of $\boldsymbol{a}$ and $\boldsymbol{b}$ is the m.p. action $\boldsymbol{a} \times \boldsymbol{b} = \Gamma \curvearrowright^{a \times b} (X \times Y, \mu \times \nu)$ where $\gamma^{a \times b}(x, y) = (\gamma^a x, \gamma^b y)$ for each $\gamma \in \Gamma$ and $(x, y) \in X \times Y$.

Bernoulli shifts. Let $\Gamma \times T \to T$, $(\gamma, t) \mapsto \gamma \cdot t$ be an action of Γ on a countable set T. The *generalized Bernoulli shift* corresponding to this action is the system $\boldsymbol{s}_{\Gamma, T} = \Gamma \curvearrowright^s ([0, 1]^T, \lambda^T)$, where λ is Lebesgue measure and where the action s is given by $(\gamma^s f)(t) = f(\gamma^{-1} \cdot t)$ for $\gamma \in \Gamma$, $f \in [0, 1]^T$, $t \in T$. We write $\boldsymbol{s}_\Gamma$ for $\boldsymbol{s}_{\Gamma, \Gamma}$ when the action of Γ on itself is by left translation. The system $\boldsymbol{s}_\Gamma$ is called the *Bernoulli shift* of Γ.

The trivial system. We call $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ *trivial* if μ is a point mass. Otherwise, $\boldsymbol{a}$ is called *non-trivial.* Up to isomorphism, each group Γ has a unique trivial measure preserving action, which we denote by $\boldsymbol{i}_\Gamma$ or simply $\boldsymbol{i}$ when Γ is clear.

Identity systems. Let $\boldsymbol{\iota}_{\Gamma, \mu} = \Gamma \curvearrowright^\iota (X, \mu)$ denote the *identity system* of Γ on (X, μ) given by $\gamma^\iota = \mathrm{id}_X$ for all $\gamma \in \Gamma$. We write $\boldsymbol{\iota}_\mu$ when Γ is clear. Thus if μ is a point mass then $\boldsymbol{\iota}_\mu \cong \boldsymbol{i}$.

Strong ergodicity. A system $\boldsymbol{a}$ is called *strongly ergodic* if it is ergodic and does not weakly contain the identity system $\boldsymbol{\iota}_{\Gamma, \lambda}$ on $([0, 1], \lambda)$.

Fixed point sets and free actions. For a subset $C \subseteq \Gamma$ we let

$$\mathrm{Fix}^b(C) = \{y \in Y : \forall \gamma \in C \ \ \gamma^b y = y\}.$$

We write $\mathrm{Fix}^b(\gamma)$ for $\mathrm{Fix}^b(\{\gamma\})$. The *kernel* of the system $\boldsymbol{b}$ is the set $\ker(\boldsymbol{b}) = \{\gamma \in \Gamma : \nu(\mathrm{Fix}^b(\gamma)) = 1\}$. It is clear that $\ker(\boldsymbol{b})$ is a normal subgroup of Γ. The system $\boldsymbol{b}$ is called *faithful* if $\ker(\boldsymbol{b}) = \{e\}$, i.e., $\nu(\mathrm{Fix}^b(\gamma)) < 1$ for each $\gamma \in \Gamma \setminus \{e\}$. The system $\boldsymbol{b}$ is called *(essentially) free* if the stabilizer of ν-almost every point is trivial, i.e., $\nu(\mathrm{Fix}^b(\gamma)) = 0$ for each $\gamma \in \Gamma \setminus \{e\}$.

2.3 Invariant random subgroups

The space of subgroups. We let $\mathrm{Sub}_\Gamma \subseteq 2^\Gamma$ denote the compact space of all subgroup of Γ and we let $c : \Gamma \times \mathrm{Sub}_\Gamma \to \mathrm{Sub}_\Gamma$ be the continuous action of Γ on Sub_Γ by conjugation.

Invariant random subgroups. An *invariant random subgroup* (IRS) of Γ is a conjugation-invariant Borel probability measures on Sub_Γ. The point mass δ_N at a normal subgroup N of Γ is an example of an invariant random subgroup. Let IRS_Γ denote the space of invariant random subgroups of Γ. We associate to each $\theta \in \mathrm{IRS}_\Gamma$ the measure preserving action $\Gamma \curvearrowright^c (\mathrm{Sub}_\Gamma, \theta)$. We also denote this system by $\boldsymbol{\theta}$.

Stabilizer distributions. Each measure preserving action $\boldsymbol{b} = \Gamma \curvearrowright^b (Y, \nu)$ of Γ gives rise to and invariant random subgroup $\theta_{\boldsymbol{b}}$ of Γ as follows. The *stabilizer* of a point $y \in Y$ under the action b is the subgroup Γ_y of Γ defined by

$$\Gamma_y = \{\gamma \in \Gamma : \gamma^b y = y\}.$$

The group Γ_y of course depends on the action b. The *stabilizer map* associated to b is the map $\mathrm{stab}_b : Y \to \mathrm{Sub}_\Gamma$ given by $\mathrm{stab}_b(y) = \Gamma_y$. The *stabilizer distribution* of $\boldsymbol{b}$, which we denote by $\theta_{\boldsymbol{b}}$ or $\mathrm{type}(\boldsymbol{b})$, is the measure $(\mathrm{stab}_b)_*\nu$ on Sub_Γ. It is clear that $\theta_{\boldsymbol{b}}$ is an invariant random subgroup of Γ. In [AGV12] it is shown that for any invariant random subgroup θ of Γ, there exists a m.p. action $\boldsymbol{b}$ of Γ such that $\theta_{\boldsymbol{b}} = \theta$. Moreover, if $\boldsymbol{\theta}$ is ergodic then $\boldsymbol{b}$ can be taken to be ergodic as well. See [CP12].

Group theoretic properties of invariant random subgroups. Let θ be an invariant random subgroup of Γ. We say that a given property $\mathcal{P}$ of subgroups of Γ holds for θ if $\mathcal{P}$ holds almost everywhere. For example, θ is called *amenable* (or *infinite index*) if θ concentrates on the amenable (respectively, infinite index) subgroups of Γ.

The trivial IRS. By the *trivial* invariant random subgroup we mean the point mass at the trivial subgroup $\{e\}$ of Γ. We write δ_e instead of $\delta_{\{e\}}$ for the trivial invariant random subgroup. An invariant random subgroup not equal to δ_e is called *non-trivial.*

Remark 2.1. We will often abuse terminology and confuse an invariant random subgroup θ with the measure preserving action $\boldsymbol{\theta} = \Gamma \curvearrowright^c (\mathrm{Sub}_\Gamma, \theta)$ it defines, stating, for example, that θ is ergodic or is weakly contained in $\boldsymbol{s}_\Gamma$ to mean that $\boldsymbol{\theta}$ is ergodic or is weakly contained in $\boldsymbol{s}_\Gamma$. When there is a potential for ambiguity we will make sure to distinguish between an invariant random subgroup and the measure preserving system to which it gives rise. We emphasise that "θ is non-trivial" will always mean that θ is not equal to the trivial IRS δ_e, whereas "$\boldsymbol{\theta}$ is non-trivial" will always mean that θ is not a point mass (at *any* subgroup).

3 Shift-minimality

3.1 Seven characterizations of shift-minimality

It will be useful to record here the main theorem of [AW11] which was already mentioned several times in the introduction.

Theorem 3.1 ([AW11])**.** *Let Γ be a countably infinite group. Then the Bernoulli shift $\boldsymbol{s}_\Gamma$ is weakly contained in every free measure preserving action of Γ.*

We let $\mathrm{Aut}(X, \mu)$ denote the Polish group of measure preserving transformations of (X, μ), and we let $A(\Gamma, X, \mu)$ denote the Polish space of measure preserving actions of Γ on the measure space (X, μ). See [Ke10] for information on these two spaces. In the following proposition, let $[\boldsymbol{a}]$ denote the weak equivalence class of a measure preserving action $\boldsymbol{a}$ of Γ. Denote by $\boldsymbol{s}_{\Gamma,2}$ the full 2-shift of Γ, i.e., the shift action of Γ on $(2^\Gamma, \rho^\Gamma)$ where we identify 2 with $\{0, 1\}$ and where $\rho(\{0\}) = \rho(\{1\}) = 1/2$.

Proposition 3.2. *Let Γ be a countable group and let (X, μ) be a standard non-atomic probability space. Then the following are equivalent.*

1. *Γ is shift-minimal, i.e., every non-trivial m.p. action weakly contained in $\boldsymbol{s}_\Gamma$ is free.*

2. *Every non-trivial m.p. action weakly contained in* $\boldsymbol{s}_{\Gamma,2}$ *is free.*

3. *Among non-trivial m.p. actions of* Γ, $[\boldsymbol{s}_{\Gamma,2}]$ *is minimal with respect to weak containment.*

4. *Either* $\Gamma = \{e\}$ *or, among non-trivial m.p. actions of* Γ, $[\boldsymbol{s}_\Gamma]$ *is minimal with respect to weak containment.*

5. *Among non-atomic m.p. actions of* Γ, $[\boldsymbol{s}_\Gamma]$ *is minimal with respect to weak containment.*

6. *The conjugation action of the Polish group* $\mathrm{Aut}(X,\mu)$ *on the Polish space* $A_{\boldsymbol{s}}(\Gamma,X,\mu) = \{\boldsymbol{a} \in A(\Gamma,X,\mu) : \boldsymbol{a} \prec \boldsymbol{s}_\Gamma\}$ *is minimal, i.e., every orbit is dense.*

7. *For some (equivalently: every) non-principal ultrafilter* $\mathcal{U}$ *on the the natural numbers* $\mathbb{N}$, *every non-trivial factor of the ultrapower* $(\boldsymbol{s}_\Gamma)_{\mathcal{U}}$ *is free.*

Proof. The equivalence (7)⇔(1) follows from [CKT-D12, Theorem 1]. For the remaining equivalences, first note that if Γ is a finite group then $\boldsymbol{s}_\Gamma$ factors onto $\boldsymbol{\iota}_\mu$, so if $\Gamma \neq \{e\}$ then Γ does not satisfy (1), (4), (5) or (6). In addition, for $\Gamma \neq \{e\}$ finite, $\boldsymbol{s}_{\Gamma,2}$ factors onto a non-trivial identity system, which shows that Γ does not satisfy (2) or (3) either. This shows that the trivial group $\Gamma = \{e\}$ is the only finite group that satisfies any of the properties (1)-(6), and it is clear the trivial group satisfies all of these properties. We may therefore assume for the rest of the proof that Γ is infinite.

(1)⇒(2): This implication is clear since $\boldsymbol{s}_{\Gamma,2}$ is a factor of $\boldsymbol{s}_\Gamma$.

(2)⇒(3): Suppose that (2) holds. By hypothesis any $\boldsymbol{a}$ weakly contained in $\boldsymbol{s}_{\Gamma,2}$ is free and thus weakly contains $\boldsymbol{s}_\Gamma$ by Theorem 3.1. (3) follows since $\boldsymbol{s}_{\Gamma 2}$ is a factor of $\boldsymbol{s}_\Gamma$.

(3)⇒(4): Since we are assuming Γ is infinite, Theorem 3.1 implies $[\boldsymbol{s}_\Gamma] = [\boldsymbol{s}_{\Gamma,2}]$, and this implication follows. (4)⇒(5) is clear.

(5)⇒(6): Suppose (5) holds. By [Ke10, Proposition 10.1] the $\mathrm{Aut}(X,\mu)$-orbit closure of any $\boldsymbol{a} \in A(\Gamma,X,\mu)$ is equal to $\{\boldsymbol{b} \in A(\Gamma,X,\mu) : \boldsymbol{b} \prec \boldsymbol{a}\}$. Thus, if $\boldsymbol{a}$ is weakly equivalent to $\boldsymbol{s}_\Gamma$, then the orbit of $\boldsymbol{a}$ is dense in $A_{\boldsymbol{s}}(\Gamma,X,\mu)$. Since $[\boldsymbol{s}_\Gamma]$ is minimal with respect to weak containment, every element of $A_{\boldsymbol{s}}(\Gamma,X,\mu)$ is weakly equivalent to $\boldsymbol{s}_\Gamma$, so has dense orbit in $A_{\boldsymbol{s}}(\Gamma,X,\mu)$. Thus, the action $\mathrm{Aut}(X,\mu) \curvearrowright A_{\boldsymbol{s}}(\Gamma,X,\mu)$ is minimal.

(6)⇒(1): Suppose that every $\boldsymbol{a} \in A_{\boldsymbol{s}}(\Gamma,X,\mu)$ has dense orbit. If $\boldsymbol{\iota}_\mu \in A_{\boldsymbol{s}}(\Gamma,X,\mu)$ then, since $\boldsymbol{\iota}_\mu$ is a fixed point for the $\mathrm{Aut}(X,\mu)$ action, $\boldsymbol{\iota}_\mu = \boldsymbol{s}_\Gamma$ and thus $\Gamma = \{e\}$. Otherwise, if $\boldsymbol{\iota}_\mu \not\prec \boldsymbol{s}_\Gamma$ then the system $\boldsymbol{s}_\Gamma$ is strongly ergodic and the group Γ is therefore non-amenable. Let $\boldsymbol{b} = \Gamma \curvearrowright^{b} (Y,\nu)$ be any non-trivial m.p. action of Γ weakly contained in $\boldsymbol{s}_\Gamma$. Then $\boldsymbol{b} \times \boldsymbol{b}$ is weakly contained in $\boldsymbol{s}_\Gamma \times \boldsymbol{s}_\Gamma \cong \boldsymbol{s}_\Gamma$ and therefore $\boldsymbol{b} \times \boldsymbol{b}$ is strongly ergodic since strong ergodicity is downward closed under weak containment (see e.g., [CKT-D12, Proposition 5.6]). In particular $\boldsymbol{b} \times \boldsymbol{b}$ is ergodic and it follows that the probability space (Y,ν) is non-atomic. The action $\boldsymbol{b}$ is then isomorphic to some action $\boldsymbol{a}$ on the non-atomic space (X,μ), and $\boldsymbol{a} \in A_{\boldsymbol{s}}(\Gamma,X,\mu)$ since $\boldsymbol{b} \prec \boldsymbol{s}_\Gamma$. By hypothesis $\boldsymbol{a}$ has dense orbit in $A_{\boldsymbol{s}}(\Gamma,X,\mu)$ so that $\boldsymbol{s}_\Gamma \sim \boldsymbol{a}$ by [Ke10, Proposition 10.1] and hence $\boldsymbol{a}$ is free, and thus $\boldsymbol{b}$ is free as well. □

Two more characterizations of shift-minimality are given in terms of amenable invariant random subgroups in Theorem 3.16 below.

3.2 NA-ergodicity

Definition 3.3. Let $\boldsymbol{a}$ be a measure preserving action of a countable group Γ. We say that $\boldsymbol{a}$ is *NA-ergodic* if the restriction of $\boldsymbol{a}$ to every non-amenable subgroup of Γ is ergodic. We say that $\boldsymbol{a}$ is *strongly NA-ergodic* if the restriction of $\boldsymbol{a}$ to every non-amenable subgroup of Γ is strongly ergodic.

Example 3.4. The central example of an NA-ergodic (and in fact, strongly NA-ergodic) action is the Bernoulli shift action $\boldsymbol{s}_\Gamma$; if $H \leqslant \Gamma$ is non-amenable then $\boldsymbol{s}_\Gamma | H \cong \boldsymbol{s}_H$ is strongly ergodic. More generally, if Γ

acts on a countable set T and the stabilizer of every $t \in T$ is amenable then the generalized Bernoulli shift $\boldsymbol{s}_T = \Gamma \curvearrowright^{s_T} ([0,1]^T, \lambda^T)$ is strongly NA-ergodic (see e.g., [KT09]).

Example 3.5. The action $\mathrm{SL}_2(\mathbb{Z}) \curvearrowright (\mathbb{T}^2, \lambda^2)$ by matrix multiplication, where λ^2 is Haar measure on $\mathbb{T}^2$, is another example of a strongly NA-ergodic action. A proof of this is given in [Ke07, 5.(B)].

Example 3.6. I would like to thank L. Bowen for bringing my attention to this example. Let Γ be a countable group and let f be an element of the integral group ring $\mathbb{Z}\Gamma$. The left translation action of Γ on the discrete abelian group $\mathbb{Z}\Gamma/\mathbb{Z}\Gamma f$ is by automorphisms, and this induces an action of Γ by automorphisms on the dual group $\widehat{\mathbb{Z}\Gamma/\mathbb{Z}\Gamma f}$, which is a compact metrizable abelian group so that this action preserves normalized Haar measure μ_f. Bowen has shown that if the function f has an inverse in $\ell^1(\Gamma)$ then the system $\Gamma \curvearrowright (\widehat{\mathbb{Z}\Gamma/\mathbb{Z}\Gamma f}, \mu_f)$ is weakly contained in $\boldsymbol{s}_\Gamma$ and is therefore strongly NA-ergodic by Proposition 3.10 ([Bo11a, §5]; note that the hypothesis that Γ is residually finite is not used in that section so that this holds for arbitrary countable groups Γ).

Remark 3.7. The actions from Examples 3.4, 3.5, and 3.6 share a common property: they are *tempered* in the sense of [Ke07]. A measure preserving action $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ is called *tempered* if the Koopman representation $\kappa_0^{\boldsymbol{a}}$ on $L_0^2(X,\mu) = L^2(X,\mu) \ominus \mathbb{C}1_X$ is weakly contained in the regular representation λ_Γ of Γ. Any tempered action $\boldsymbol{a}$ of a non-amenable group Γ has *stable spectral gap* in the sense of [Pop08] (this means $\kappa_0^{\boldsymbol{a}} \otimes \kappa_0^{\boldsymbol{a}}$ does not weakly contain the trivial representation), and this implies in turn that the product action $\boldsymbol{a} \times \boldsymbol{b}$ is strongly ergodic relative to $\boldsymbol{b}$ for every measure preserving action $\boldsymbol{b}$ of Γ (see [Io10]). In particular (taking $\boldsymbol{b} = \boldsymbol{i}_\Gamma$) a tempered action $\boldsymbol{a}$ of a non-amenable group is itself strongly ergodic. Since the restriction of a tempered action to a subgroup is still tempered it follows that every tempered action is strongly NA-ergodic. In [Ke07] it is shown that the converse holds for any action on a compact Polish group G by automorphisms (such an action necessarily preserves Haar measure μ_G):

Theorem 3.8 (Theorem 4.6 of [Ke07]). *Let Γ be a countably infinite group acting by automorphisms on a compact Polish group G. Let $\hat{G}$ denote the (countable) set of all isomorphism classes of irreducible unitary representations of G and let $\hat{G}_0 = \hat{G} \setminus \{\hat{1}_G\}$. Then the following are equivalent:*

1. *The action $\Gamma \curvearrowright (G, \mu_G)$ is tempered;*
2. *Every stabilizer of the associated action of Γ on $\hat{G}_0$ is amenable.*
3. *The action $\Gamma \curvearrowright (G, \mu_G)$ is NA-ergodic.*
4. *The action $\Gamma \curvearrowright (G, \mu_G)$ is strongly NA-ergodic.*

Condition (2) of Theorem 3.8 should be compared with part (ii) of Lemma 3.11 below, although Lemma 3.11 deals with general NA-ergodic actions. It follows from [Ke10, Proposition 10.5] that any measure preserving action weakly contained in $\boldsymbol{s}_\Gamma$ is tempered. I do not know however whether the converse holds, although Example 3.6 and Theorem 3.8 suggest that this may be the case for actions by automorphisms on compact Polish groups.

Question 3.9. Let Γ be a countable group acting by automorphisms on a compact Polish group G and assume the action is tempered. Does it follow that the action is weakly contained in $\boldsymbol{s}_\Gamma$? As a special case, is it true that the action $\mathrm{SL}_2(\mathbb{Z}) \curvearrowright (\mathbb{T}^2, \lambda^2)$ is weakly contained in $\boldsymbol{s}_{\mathrm{SL}_2(\mathbb{Z})}$?

We now establish some properties of general NA-ergodic actions.

Proposition 3.10. *Any factor of an NA-ergodic action is NA-ergodic. Any action weakly contained in a strongly NA-ergodic action is strongly NA-ergodic.*

Proof. The first statement is clear and the second is a consequence of strong ergodicity being downward closed under weak containment (see [CKT-D12, Proposition 5.6]). ☐

Part (ii) of the following lemma is one of the key facts about NA-ergodicity.

Lemma 3.11. *Let* $\boldsymbol{b} = \Gamma \curvearrowright^b (Y, \nu)$ *be any non-trivial NA-ergodic action of a countable group* Γ*.*

(i) Suppose that $C \subseteq \Gamma$ *is a subset of* Γ *such that* $\nu(\{y \in Y : C \subseteq \Gamma_y\}) > 0$*. Then the subgroup* $\langle C \rangle$ *generated by* C *is amenable.*

(ii) The stabilizer Γ_y *of* ν*-almost every* $y \in Y$ *is amenable.*

Proof. We begin with part (i). The hypothesis tells us that $\nu(\mathrm{Fix}^b(C)) > 0$. Since ν is not a point mass there is some $B \subseteq \mathrm{Fix}^b(C)$ with $0 < \nu(B) < 1$. Then B witnesses that $\boldsymbol{b} \restriction \langle C \rangle$ is not ergodic, so $\langle C \rangle$ is amenable by NA-ergodicity of $\boldsymbol{b}$.

For (ii), let $\mathcal{F}$ denote the collection of finite subsets F of Γ such that $\langle F \rangle$ is non-amenable and let $\mathrm{NA} = \{y \in Y : \Gamma_y \text{ is non-amenable}\}$. Then

$$\mathrm{NA} = \bigcup_{F \in \mathcal{F}} \{y \in Y : F \subseteq \Gamma_y\}.$$

By part (i), $\nu(\{y \in Y : F \subseteq \Gamma_y\}) = 0$ for each $F \in \mathcal{F}$. Since $\mathcal{F}$ is countable it follows that $\nu(\mathrm{NA}) = 0$. ☐[Lemma]

The function $N : \mathrm{Sub}_\Gamma \to \mathrm{Sub}_\Gamma$ sending a subgroup $H \leqslant \Gamma$ to its normalizer $N(H)$ in Γ is equivariant for the conjugation action $\Gamma \curvearrowright^c \mathrm{Sub}_\Gamma$. In [Ve12, §2.4] Vershik examines the following transfinite iterations of this function.

Definition 3.12. Define $N^\alpha : \mathrm{Sub}_\Gamma \to \mathrm{Sub}_\Gamma$ by transfinite induction on ordinals α as follows.

$$\begin{aligned} N^0(H) &= H, \\ N^{\alpha+1}(H) &= N(N^\alpha(H)) \text{ is the normalizer of } N^\alpha(H) \\ N^\lambda(H) &= \bigcup_{\alpha<\lambda} N^\alpha(H) \text{ when } \lambda \text{ is a limit ordinal.} \end{aligned}$$

Each N^α is equivariant with respect to conjugation. For each H the sets $H, N(H), \dots, N^\alpha(H), N^{\alpha+1}(H), \dots$ form an increasing ordinal-indexed sequence of subsets of Γ. The least ordinal α_H such that $N^{\alpha_H+1}(H) = N^{\alpha_H}(H)$ is therefore countable. If $\theta \in \mathrm{IRS}_\Gamma$ then we let $\theta^\alpha = (N^\alpha)_*\theta$ for each countable ordinal $\alpha < \omega_1$. The net $\{\theta^\alpha\}_{\alpha<\omega_1}$ is increasing in the sense of [CP12, §3.5] (see also the paragraphs preceding Theorem A.15 below), so by [CP12, Theorem 3.12] there is a weak*-limit θ^∞ such that $\theta^\alpha \leqslant \theta^\infty$ for all α. Since IRS_Γ is a second-countable topological space there is a countable ordinal α such that $\theta^\beta = \theta^\infty$ for all $\beta \geqslant \alpha$. Thus $N_*\theta^\infty = \theta^\infty$, and it follows from [Ve12, Proposition 4] that θ^∞ concentrates on the self-normalizing subgroups of Γ.

Theorem 3.13. *Let* $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \nu)$ *be a non-trivial measure preserving action of the countable group* Γ*. Suppose that* $\boldsymbol{a}$ *is NA-ergodic. Then the stabilizer* Γ_x *of* μ*-almost every* $x \in X$ *is amenable. In addition, at least one of the following is true:*

(1) There exists a normal amenable subgroup $N \lhd \Gamma$ *such that the stabilizer of* μ*-almost every* $x \in X$ *is contained in* N*.*

(2) $\theta^\infty_{\boldsymbol{a}}$ *is a non-atomic, self-normalizing, infinitely generated amenable invariant random subgroup, where* $\theta_{\boldsymbol{a}}$ *denotes the stabilizer distribution of* $\boldsymbol{a}$*.*

Proof. Let $\theta = \theta_{\boldsymbol{a}}$. It is enough to show that either (1) or (2) is true. We may assume that Γ is non-amenable. There are two cases to consider.

Case 1: There is some ordinal α such that the measure θ^{α} has an atom. Let α_0 be the least such ordinal. Then $\boldsymbol{\theta}^{\alpha_0}$ is NA-ergodic, being a factor of $\boldsymbol{a}$, and thus the restriction of $\boldsymbol{\theta}^{\alpha_0}$ to every finite index subgroup of Γ is ergodic since Γ is non-amenable. Thus, θ^{α_0} having an atom implies that it is a point mass, so let $N \leqslant \Gamma$ be such that $\theta^{\alpha_0} = \delta_N$. Then N is a normal subgroup of Γ and we show that N is amenable so that alternative (1) holds in this case. By definition of α_0, $\boldsymbol{a}$ and each $\boldsymbol{\theta}^{\alpha}$ for $\alpha < \alpha_0$ are non-trivial NA-ergodic actions. Lemma 3.11 then implies that the invariant random subgroups $\mathrm{type}(\boldsymbol{a}) = \theta^0$ and $\mathrm{type}(\boldsymbol{\theta}^{\alpha}) = \theta^{\alpha+1}$, for $\alpha < \alpha_0$, all concentrate on the amenable subgroups of Γ. If $\alpha_0 = 0$ or if α_0 is a successor ordinal then we see immediately that N is amenable. If α_0 is a limit ordinal then N is an increasing union of amenable groups and so is amenable in this case as well.

Case 2: The other possibility is that θ^{∞} has no atoms. Thus $\boldsymbol{\theta}^{\infty}$ is a non-trivial NA-ergodic action with $\mathrm{type}(\boldsymbol{\theta}^{\infty}) = N_*\theta^{\infty} = \theta^{\infty}$. This implies that θ^{∞} is amenable by Lemma 3.11. Since θ^{∞} is non-atomic and there are only countably many finitely generated subgroups of Γ, θ^{∞} must concentrate on the infinitely-generated subgroups. This shows that (2) holds. ☐

3.3 Amenable invariant random subgroups

We record a corollary of Theorem 3.13 which will be used in the proof of our final characterization of shift-minimality.

Corollary 3.14. *Any group Γ that is not shift-minimal either has a non-trivial normal amenable subgroup N, or has a non-atomic, self-normalizing, infinitely-generated, amenable invariant random subgroup θ such that the action $\boldsymbol{\theta} = \Gamma \curvearrowright^{c} (\mathrm{Sub}_{\Gamma}, \theta)$ is weakly contained in $\boldsymbol{s}_{\Gamma}$.*

Proof. Let Γ be a group that is not shift-minimal so that there exists some non-trivial $\boldsymbol{a}$ weakly contained in $\boldsymbol{s}_{\Gamma}$ which is not free. The action $\boldsymbol{a}$ is strongly NA-ergodic by 3.4 and 3.10, so $\boldsymbol{a}$ satisfies the hypotheses of Theorem 3.13. If (1) of Theorem 3.13 holds, say with witnessing normal amenable subgroup $N \leqslant \Gamma$, then N is non-trivial since $\boldsymbol{a}$ is non-free. If alternative (2) of Theorem 3.13 holds then taking $\theta = \theta^{\infty}_{\boldsymbol{a}}$ works. ☐

We also need

Proposition 3.15. *If Γ is shift-minimal then Γ has no non-trivial normal amenable subgroups.*

Proof. Suppose that Γ has a non-trivial normal amenable subgroup N. Amenability implies that $\boldsymbol{\iota}_N \prec \boldsymbol{s}_N$. Then since co-inducing preserves weak containment we have

$$\boldsymbol{s}_{\Gamma,\Gamma/N} \prec \mathrm{CInd}_N^{\Gamma}(\boldsymbol{\iota}_N) \prec \mathrm{CInd}_N^{\Gamma}(\boldsymbol{s}_N) \cong \boldsymbol{s}_{\Gamma}$$

which shows that $\boldsymbol{s}_{\Gamma,\Gamma/N} \prec \boldsymbol{s}_{\Gamma}$. The action $\boldsymbol{s}_{\Gamma,\Gamma/N}$ is not free since $N \subseteq \ker(\boldsymbol{s}_{\Gamma,\Gamma/N})$. This shows that Γ is not shift-minimal. ☐

The following immediately yields Theorem 1.3 from the introduction.

Theorem 3.16. *The following are equivalent for a countable group Γ:*

(1) Γ is not shift-minimal.

(2) There exists a non-trivial amenable invariant random subgroup θ of Γ that is weakly contained in $\boldsymbol{s}_{\Gamma}$.

(3) Either AR_{Γ} is finite and non-trivial, or there exists an infinite amenable invariant random subgroup θ of Γ that is weakly contained in $\boldsymbol{s}_{\Gamma}$.

Proof. (1)⇒(3): Suppose that Γ is not shift-minimal. If the second alternative of Corollary 3.14 holds then we are done. Otherwise, the first alternative holds and so AR_Γ is non-trivial. If AR_Γ is finite then (3) is immediate, and if AR_Γ is infinite then the point-mass at AR_Γ shows that (3) holds.

(3)⇒(2) is clear. Now let θ be as in (2) and we will show that Γ is not shift-minimal. If θ is a point mass, say at $H \in \mathrm{Sub}(\Gamma)$, then H is normal and by hypothesis H is non-trivial and amenable so (1) then follows from Proposition 3.15. If θ is not a point mass then $\Gamma \curvearrowright^c (\mathrm{Sub}_\Gamma, \theta)$ is a non-trivial and non-free measure preserving action of Γ that is weakly contained in $\boldsymbol{s}_\Gamma$. This action then witnesses that Γ is not shift-minimal. □

Any group with no non-trivial normal amenable subgroups is ICC (see [Ha07, Appendix J] for a proof), so Proposition 3.15 also shows

Proposition 3.17. *Shift-minimal groups are* ICC.

4 Permanence properties

This section examines various circumstances in which shift-minimality is preserved. §4.1 establishes a lemma which will be used to show that, in many cases, shift-minimality passes to finite index subgroups.

4.1 Invariant random subgroups with trivial intersection

For each invariant random subgroup θ of Δ define the set

$$P_\theta = \{\delta \in \Delta : \theta(\{H : \delta \in H\}) > 0\}.$$

We say that two invariant random subgroups θ and ρ *intersect trivially* if $P_\theta \cap P_\rho = \{e\}$. This notion comes from looking at freeness of a product action.

Lemma 4.1. *If $\boldsymbol{a} = \Delta \curvearrowright^a (X, \mu)$ and $\boldsymbol{b} = \Delta \curvearrowright^b (Y, \nu)$ are measure preserving actions of Δ then $\boldsymbol{a} \times \boldsymbol{b}$ is free if and only if $\theta_{\boldsymbol{a}}$ and $\theta_{\boldsymbol{b}}$ intersect trivially.*

Proof. For each $\delta \in \Delta$ we have $\mathrm{Fix}^{a\times b}(\delta) = \mathrm{Fix}^a(\delta) \times \mathrm{Fix}^b(\delta)$, and this set is $(\mu \times \nu)$-null if and only if either $\mathrm{Fix}^a(\delta)$ is μ-null or $\mathrm{Fix}^b(\delta)$ is ν-null. The lemma easily follows. □

It is a straightforward group theoretic fact that if L and K are normal subgroups of Δ which intersect trivially then they commute. This generalizes to invariant random subgroups as follows.

Lemma 4.2. *Let Δ be a countable group. Let $\theta, \rho \in \mathrm{IRS}_\Delta$ and suppose that θ and ρ intersect trivially. Suppose L and K are subgroups of Δ satisfying*

$$\theta(\{H \in \mathrm{Sub}_\Delta : L \leqslant H\}) > \tfrac{1}{m}$$
$$\rho(\{H \in \mathrm{Sub}_\Delta : K \leqslant H\}) > \tfrac{1}{n}$$

for some $n, m \in \mathbb{N}$. Then there exist commuting subgroups $L_0 \leqslant L$ and $K_0 \leqslant K$ with $[L : L_0] < n$ and $[K : K_0] < m$.

Proof. Define the sets

$$Q_L = \{l \in L : \langle lKl^{-1} \cup K\rangle \subseteq P_\rho\}$$
$$Q_K = \{k \in K : \langle kLk^{-1} \cup L\rangle \subseteq P_\theta\}.$$

If $l \in Q_L$ then for any $k \in K$ we have $lkl^{-1}k^{-1} \in \langle lKl^{-1} \cup K \rangle \subseteq P_\rho$. Similarly, if $k \in Q_K$ then for any $l \in L$ we have $lkl^{-1}k^{-1} \in \langle kLk^{-1} \cup L \rangle \subseteq P_\theta$. Thus, if $l \in Q_L$ and $k \in Q_K$ then $lkl^{-1}k^{-1} \in P_\rho \cap P_\theta = \{e\}$ and so l and k commute. It follows that the groups $L_0 = \langle Q_L \rangle \leqslant L$ and $K_0 = \langle Q_K \rangle \leqslant K$ commute.

Suppose for contradiction that $[L : L_0] \geqslant n$ and let $l_0, \dots, l_{n-1}$ be elements of distinct left cosets of L_0 in L, with $l_0 = e$. For each $i < n$ let $A_i = \{H \in \mathrm{Sub}_\Delta \, : \, l_i K l_i^{-1} \leqslant H\}$ so that $\rho(A_i) = \rho(l_i^c \cdot A_0) = \rho(A_0) > \frac{1}{n}$ by hypothesis. There must be some $0 \leqslant i < j < n$ with $\rho(A_i \cap A_j) > 0$. Let $l = l_j^{-1} l_i$. Then $\rho(l^c \cdot A_0 \cap A_0) = \rho(A_i \cap A_j) > 0$ and $l^c \cdot A_0 \cap A_0$ consists of those $H \in \mathrm{Sub}_\Delta$ such that $lKl^{-1} \cup K \leqslant H$. This shows that $\langle lKl^{-1} \cup K \rangle \subseteq P_\rho$ and thus $l \in Q_L \subseteq L_0$. But this contradicts that $l = l_j^{-1} l_i$ and $l_i L_0 \neq l_j L_0$. Therefore $[L : L_0] < n$. Similarly, $[K : K_0] < m$. ☐[Lemma 4.2]

Theorem 4.3. *Let $\theta, \rho \in \mathrm{IRS}_\Delta$, $L, K \leqslant \Delta$, and $n, m \in \mathbb{N}$ be as in Lemma 4.2, and assume in addition that L and K are finitely generated. Then there exist commuting subgroups N_L and N_K, both normal in Δ, with $[L : L \cap N_L] < \infty$ and $[K : K \cap N_K] < \infty$.*

Proof. For a subgroup $H \leqslant \Delta$ and $i \in \mathbb{N}$ let $H(i)$ be the intersection of all subgroups of H of index strictly less than i. Then $L(n)$ is finite index in L, and $K(m)$ is finite index in K, since L and K are finitely generated. By Lemma 4.2 $L(n)$ and $K(m)$ commute. For any $\gamma, \delta \in \Delta$ the groups $\gamma L \gamma^{-1}$ and $\delta K \delta^{-1}$ satisfy the hypotheses of Lemma 4.2 hence the groups $(\gamma L \gamma^{-1})(n) = \gamma L(n) \gamma^{-1}$ and $(\delta K \delta^{-1})(m) = \delta K(m) \delta^{-1}$ commute. It follows that the normal subgroups $N_L = \langle \bigcup_{\delta \in \Delta} \delta L(n) \delta^{-1} \rangle$ and $N_K = \langle \bigcup_{\delta \in \Delta} \delta K(m) \delta^{-1} \rangle$ satisfy the conclusion of the theorem. ☐

4.2 Finite index subgroups

The following is an analogue of a theorem of [Be91], and its proof is essentially the same as [BH00, Proposition 6].

Proposition 4.4. *Let $\boldsymbol{a}$ be a measure preserving action of a countable group Γ and let N be a normal subgroup of Γ. If the restriction $\boldsymbol{a} \restriction N$ of $\boldsymbol{a}$ to N is free then $\mu(\mathrm{Fix}^a(\gamma)) = 0$ for any $\gamma \in \Gamma$ satisfying*

$$|\{h \gamma h^{-1} \, : \, h \in N\}| = \infty. \tag{4.1}$$

Thus, if (4.1) *holds for all $\gamma \notin N$ then a m.p. action of Γ is free if and only if its restriction to N is free.*

For example, it is shown in [Be91] that (4.1) holds for all $\gamma \notin N$ whenever $C_\Gamma(N) = \{e\}$ and N is ICC.

Proof of Proposition 4.4. Suppose $\gamma \in \Gamma \setminus \{e\}$ is such that $\mu(\mathrm{Fix}^a(\gamma)) > 0$ and $\{h \gamma h^{-1} \, : \, h \in N\}$ is infinite. It suffices to show that $\boldsymbol{a} \restriction N$ is not free. The Poincaré recurrence theorem implies that there exist $h_0, h_1 \in N$ with $h_0 \gamma h_0^{-1} \neq h_1 \gamma h_1^{-1}$ and $\mu(h_0^a \cdot \mathrm{Fix}^a(\gamma) \cap h_1^a \cdot \mathrm{Fix}^a(\gamma)) > 0$. Let $h = h_1^{-1} h_0$ so that $h \in N$ and $h \gamma h^{-1} \neq \gamma$. Since $\mathrm{Fix}^a(\gamma) = \mathrm{Fix}^a(\gamma^{-1})$ we have

$$h^a \cdot \mathrm{Fix}^a(\gamma) \cap \mathrm{Fix}^a(\gamma) = \mathrm{Fix}^a(h \gamma h^{-1}) \cap \mathrm{Fix}^a(\gamma^{-1}) \subseteq \mathrm{Fix}^a(\gamma^{-1} h \gamma h^{-1}),$$

which implies $\mu(\mathrm{Fix}^a(\gamma^{-1} h \gamma h^{-1})) > 0$. This shows $\boldsymbol{a} \restriction N$ is not free since $e \neq \gamma^{-1}(h \gamma h^{-1}) = (\gamma^{-1} h \gamma) h^{-1} \in N$ by our choice of h. ☐

Proposition 4.5. *Let K be a finite index subgroup of a countable ICC group Γ, and let $\boldsymbol{a}$ be a measure preserving action of Γ. If $\boldsymbol{a} \restriction K$ is free, then $\boldsymbol{a}$ is free.*

Proof. Let $N = \bigcap_{\gamma \in \Gamma} \gamma K \gamma^{-1}$ be the normal core of K in Γ. Then N is a normal finite index subgroup of Γ. Since Γ is ICC, the group $C_\Gamma(\gamma)$ is infinite index in Γ for any $\gamma \in \Gamma$, hence $C_\Gamma(\gamma) \cap N$ is infinite index in N. In particular $\{h \gamma h^{-1} \, : \, h \in N\}$ is infinite. If $\boldsymbol{a}$ is any m.p. action of Γ whose restriction to K is free, then the restriction of $\boldsymbol{a}$ to N is free, so by Proposition 4.4, $\boldsymbol{a}$ is free. ☐

Proposition 4.5 can be used to characterize exactly when shift-minimality of Γ may be deduced from shift-minimality of one of its finite index subgroups.

Proposition 4.6. *Let* K *be a finite index subgroup of the countable group* Γ*. Suppose that* K *is shift-minimal. Then the following are equivalent.*

1. Γ *is shift-minimal.*
2. Γ *is* ICC.
3. Γ *has no non-trivial finite normal subgroups.*
4. $C_\Gamma(N) = \{e\}$ *where* $N = \bigcap_{\gamma\in\Gamma} \gamma K \gamma^{-1}$.

Proof. Since K is shift-minimal, it is also ICC by Proposition 3.15. The equivalence of (2), (3), and (4) then follows from [Pré12, Proposition 6.3]. It remains to show that (2)⇒(1). Suppose that Γ is ICC and that $\boldsymbol{a} \prec \boldsymbol{s}_\Gamma$ is non-trivial. Then $\boldsymbol{a} \upharpoonright K \prec \boldsymbol{s}_K$, so $\boldsymbol{a} \upharpoonright K$ is free by shift-minimality of K, and therefore $\boldsymbol{a}$ itself is free by Proposition 4.5. ☐

Proposition 4.6 shows that, except for the obvious counterexamples, shift-minimality is inherited from a finite index subgroup. It seems likely that, conversely, shift-minimality passes from a group to each of its finite index subgroups. By Proposition 4.6 to show this it would be enough to show that shift-minimality passes to finite index *normal* subgroups (see the discussion following Question 7.11 in §7). Theorem 4.3 can be used to give a partial confirmation of this. Recall that a group is *locally finite* if each of its finitely generated subgroups is finite.

Theorem 4.7. *Let* N *be a normal finite index subgroup of a shift-minimal group* Γ*. Suppose that* N *has no infinite locally finite invariant random subgroups that are weakly contained in* $\boldsymbol{s}_N$*. Then* N *is shift-minimal.*

Corollary 4.8. *Let* Γ *be a shift-minimal group. Then every finite index subgroup of* Γ *which is torsion-free is shift-minimal.*

Proof of Corollary 4.8. Let K be a torsion-free finite index subgroup of Γ. Note that K is ICC since the ICC property passes to finite index subgroups. The group $N := \bigcap_{\gamma\in\Gamma} \gamma K \gamma^{-1}$ is finite index in Γ and torsion-free, and it is moreover normal in Γ. By Theorem 4.7, N is shift-minimal, whence K is shift-minimal by Proposition 4.6. ☐

Theorem 4.7 will follow from:

Theorem 4.9. *Let* Δ *be a countable group with* $\mathrm{AR}_\Delta = \{e\}$*. Let* θ *and* ρ *be invariant random subgroups of* Δ *which are not locally finite. Suppose that* $\boldsymbol{\rho}$ *is* NA*-ergodic. Then* θ *and* ρ *have non-trivial intersection.*

We first show how to deduce 4.7 from 4.9.

Proof of Theorem 4.7 from Theorem 4.9. Let $\boldsymbol{a} = N \curvearrowright^a (X,\mu)$ be a non-trivial m.p. action of N weakly contained in $\boldsymbol{s}_N$. We will show that $\boldsymbol{a}$ is free.

The coinduced action $\boldsymbol{c} = \mathrm{CInd}_N^\Gamma(\boldsymbol{a})$ is weakly contained in $\boldsymbol{s}_\Gamma$, so $\boldsymbol{c}$ is free by shift-minimality of Γ. Let $T = \{t_0, \ldots, t_{n-1}\}$ be a transversal for the left cosets of N in Γ. Then $\boldsymbol{c} \upharpoonright N \cong \prod_{0\leqslant i<n} \boldsymbol{a}^{t_i}$ where for $\boldsymbol{b} \in A(N, X, \mu)$, $\boldsymbol{b}^t \in A(N, X, \mu)$ is given by $k^{b^t} = (t^{-1}kt)^b$ for each $k \in N$, $t \in T$ [Ke10, 10.**(G)**]. Observe that $\theta_{\boldsymbol{a}^t} = (\varphi_t)_*\theta_{\boldsymbol{a}}$ where $\varphi_t : \mathrm{Sub}_N \to \mathrm{Sub}_N$ is the conjugation map $H \mapsto tHt^{-1}$. In particular, for each $t \in T$, $\boldsymbol{a}^t$ is free if and only if $\boldsymbol{a}$ is free. It is easy to see that $(\boldsymbol{s}_N)^t \cong \boldsymbol{s}_N$ for each $t \in T$, so it follows

that $\boldsymbol{c} \restriction N \cong \prod_{0 \leqslant i < n} \boldsymbol{a}^{t_i} \prec \boldsymbol{s}_N$. For each $j < n$ let $\boldsymbol{c}_j = \prod_{j \leqslant i < n} \boldsymbol{a}^{t_i}$. We will show that $\boldsymbol{c}_j$ is free for all $0 \leqslant j < n$, which will finish the proof since this will show that $\boldsymbol{c}_{n-1} = \boldsymbol{a}^{t_{n-1}}$ is free, whence $\boldsymbol{a}$ is free.

We know that $\boldsymbol{c}_0 = \boldsymbol{c} \restriction N$ is free. Assume for induction that $\boldsymbol{c}_{j-1}$ is free (where $j \geqslant 1$ is less than n) and we will show that $\boldsymbol{c}_j$ is free. Note the following:

(i) $\theta_{\boldsymbol{a}^{t_{j-1}}}$ and $\theta_{\boldsymbol{c}_j}$ intersect trivially. This follows from Lemma 4.1 because $\boldsymbol{c}_{j-1} = \boldsymbol{a}^{t_{j-1}} \times \boldsymbol{c}_j$ is free.

(ii) Both $\boldsymbol{\theta}_{\boldsymbol{a}^{t_{j-1}}}$ and $\boldsymbol{\theta}_{\boldsymbol{c}_j}$ are NA-ergodic, since they are both weakly contained in $\boldsymbol{s}_N$.

(iii) $\mathrm{AR}_N = \{e\}$. This is because Γ is shift-minimal, so that $\mathrm{AR}_\Gamma = \{e\}$ by Proposition 3.15, and N is normal in Γ so apply Proposition B.1.

Theorem 4.9 along with (i), (ii), and (iii) imply that either $\theta_{\boldsymbol{a}^{t_{j-1}}}$ or $\theta_{\boldsymbol{c}_j}$ is locally finite. But N has no infinite locally finite invariant random subgroups weakly contained in $\boldsymbol{s}_N$ by hypothesis, and since $\mathrm{AR}_N = \{e\}$, N actually has no *non-trivial* locally finite invariant random subgroups weakly contained in $\boldsymbol{s}_N$. It follows that either $\theta_{\boldsymbol{a}^{t_{j-1}}}$ or $\theta_{\boldsymbol{c}_j}$ is trivial. If $\theta_{\boldsymbol{c}_j}$ is trivial then $\boldsymbol{c}_j$ is free, which is what we wanted to show. If $\theta_{\boldsymbol{a}^{t_{j-1}}}$ is trivial then $\boldsymbol{a}^{t_{j-1}}$ is free, so $\boldsymbol{a}^{t_i}$ is free for all $i < n$, and therefore $\boldsymbol{c}_j$ is free all the same. ☐

Proof of Theorem 4.9. Suppose toward a contradiction that θ and ρ intersect trivially. By hypothesis θ is not locally finite, so the set of $H \in \mathrm{Sub}_\Delta$ that contain an infinite finitely generated subgroup is θ-non-null. As there are only countably many infinite finitely generated subgroups of Δ, there must be at least one – call it L – for which $\theta(\{H : L \subseteq H\}) > 0$. Similarly, there is an infinite finitely generated $K \leqslant \Delta$ with $\rho(\{H : K \leqslant H\}) > 0$. Then θ, ρ, L and K satisfy the hypotheses of Theorem 4.3 (for some n and m), so there exist normal subgroups $N_L, N_K \leqslant \Delta$ which commute, with $[L : L \cap N_L] < \infty$ and $[K : K \cap N_K] < \infty$. Since L and K are infinite, neither N_L nor N_K is trivial, and since $\mathrm{AR}_\Delta = \{e\}$, both N_L and N_K are non-amenable.

Pick some $k \neq e$ with $k \in K \cap N_K$. Since $k \in K$, the set $\{H : k \in H\}$ has positive ρ-measure, and it is N_L-invariant since N_L commutes with k. NA-ergodicity of $\boldsymbol{\rho}$ and non-amenability of N_L then imply that $\rho(\{H : k \in H\}) = 1$. On the other hand, the set

$$M_\rho = \{\delta \in \Delta : \rho(\{H : \delta \in H\}) = 1\}$$

is a normal subgroup of Δ which acts trivially under $\boldsymbol{\rho}$, so NA-ergodicity of $\boldsymbol{\rho}$ implies M_ρ is amenable, and as $\mathrm{AR}_\Delta = \{e\}$, we actually have $M_\rho = \{e\}$, which contradicts that $k \in M_\rho$. ☐

Question 7.11 below asks whether a finite index subgroup of a shift-minimal group is always shift-minimal.

4.3 Direct sums

Proposition 4.10. *Let* $(\Gamma_i)_{i \in I}$ *be a sequence of countable* ICC *groups and let* $\boldsymbol{a}$ *be a measure preserving action of* $\Gamma = \bigoplus_{i \in I} \Gamma_i$. *If* $\boldsymbol{a} \restriction \Gamma_i$ *is free for each* $i \in I$ *then* $\boldsymbol{a}$ *is free. In particular, the direct sum of shift-minimal groups is shift-minimal.*

Proof. We will show that if $\boldsymbol{a}$ is not free then $\boldsymbol{a} \restriction \Gamma_i$ is not free for some $i \in I$. We give the proof for the case of the direct sum of two ICC groups – say Γ_1 and Γ_2 – since the proof for infinitely many groups is nearly identical. Let $\Gamma = \Gamma_1 \times \Gamma_2$ and let $(\gamma, \delta) \in \Gamma$ be such that $\mu(\mathrm{Fix}^a((\gamma, \delta))) > 0$ where $(\gamma, \delta) \neq e_\Gamma$. Suppose that $\delta \neq e$ (the case where $\gamma \neq e$ is similar). Since Γ_2 is ICC we have that $C_{\Gamma_2}(\delta)$ is infinite index in Γ_2 so by Poincaré recurrence there exists $\alpha \in \Gamma_2$, $\alpha \notin C_{\Gamma_2}(\delta)$ such that

$$\mu\big((e, \alpha)^a \cdot \mathrm{Fix}^a((\gamma, \delta)) \cap \mathrm{Fix}^a((\gamma, \delta))\big) > 0.$$

Thus $\mu\big(\mathrm{Fix}^a(\langle(\gamma, \alpha\delta\alpha^{-1}), (\gamma, \delta)\rangle)\big) > 0$ and in particular $\mu\big(\mathrm{Fix}^a((e, \alpha\delta\alpha^{-1}\delta^{-1}))\big) > 0$. Our choice of α implies that $\alpha\delta\alpha^{-1}\delta^{-1} \neq e$ and so $\boldsymbol{a} \restriction \Gamma_2$ is non-free as was to be shown. ☐

4.4 Other permanence properties

Proposition 4.11. *Let $\boldsymbol{a}$ be a measure preserving action of Γ. Let N be a normal subgroup of Γ. Suppose that both N and $\mathrm{C}_\Gamma(\mathrm{N})$ are ICC. Suppose that $\boldsymbol{a} \upharpoonright \mathrm{N}$ and $\boldsymbol{a} \upharpoonright \mathrm{C}_\Gamma(\mathrm{N})$ are both free. Then $\boldsymbol{a}$ is free.*

Proof. Let $\mathrm{K} = \mathrm{C}_\Gamma(\mathrm{N})\mathrm{N}$. Then K is normal in Γ since both N and $\mathrm{C}_\Gamma(\mathrm{N})$ are normal. By hypothesis $\mathrm{C}_\Gamma(\mathrm{N}) \cap \mathrm{N} = \{e\}$ so $\mathrm{K} \cong \mathrm{C}_\Gamma(\mathrm{N}) \times \mathrm{N}$. It follows that K is ICC, being a product of ICC groups. Proposition 4.10 then implies that $\boldsymbol{a} \upharpoonright \mathrm{K}$ is free. Since $\mathrm{C}_\Gamma(\mathrm{K}) \leqslant \mathrm{C}_\Gamma(\mathrm{C}_\Gamma(\mathrm{N})) \cap \mathrm{C}_\Gamma(\mathrm{N}) = \mathrm{Z}(\mathrm{C}_\Gamma(\mathrm{N})) = \{e\}$, Proposition 4.4 implies that $\boldsymbol{a}$ is free. ☐

Definition 4.12. A subgroup H of Γ is called *almost ascendant* in Γ if there exists a well-ordered increasing sequence $\{\mathrm{H}_\alpha\}_{\alpha \leqslant \lambda}$ of subgroups of Γ, indexed by some countable ordinal λ, such that

(i) $\mathrm{H} = \mathrm{H}_0$ and $\mathrm{H}_\lambda = \Gamma$.

(ii) For each $\alpha < \lambda$, either H_α is a normal subgroup of $\mathrm{H}_{\alpha+1}$ or H_α is a finite index subgroup of $\mathrm{H}_{\alpha+1}$.

(iii) $\mathrm{H}_\beta = \bigcup_{\alpha<\beta} \mathrm{H}_\alpha$ whenever β is a limit ordinal.

We call $\{\mathrm{H}_\alpha\}_{\alpha \leqslant \lambda}$ an *almost ascendant series* for H in Γ. If H is almost ascendant in Γ and if there exists an almost ascendant series $\{\mathrm{H}_\alpha\}_{\alpha \leqslant \lambda}$ for H in Γ such that H_α is normal in $\mathrm{H}_{\alpha+1}$ for all $\alpha < \lambda$ then we say that H is *ascendant* in Γ and we call $\{\mathrm{H}_\alpha\}_{\alpha \leqslant \lambda}$ an *ascendant series* for H in Γ.

Proposition 4.13. *Let $\boldsymbol{a} = \Gamma \curvearrowright^{a} (X, \mu)$ be a measure preserving action of Γ.*

1. *Suppose that L is an almost ascendant subgroup of Γ that is ICC and satisfies $\mathrm{C}_\Gamma(\mathrm{L}) = \{e\}$. Then $\boldsymbol{a}$ is free if and only if $\boldsymbol{a} \upharpoonright \mathrm{L}$ is free. Thus, if L is shift-minimal then so is Γ.*

2. *Suppose that L is an ascendant subgroup of Γ such that $\mathrm{AR}_\mathrm{L} = \mathrm{AR}_{\mathrm{C}_\Gamma(\mathrm{L})} = \{e\}$. Then $\boldsymbol{a}$ is free if and only if both $\boldsymbol{a} \upharpoonright \mathrm{L}$ and $\boldsymbol{a} \upharpoonright \mathrm{C}_\Gamma(\mathrm{L})$ are free.*

Proof. (1): Assume that $\boldsymbol{a} \upharpoonright \mathrm{L}$ is free. Let $\{\mathrm{L}_\alpha\}_{\alpha \leqslant \lambda}$ be an almost ascendant series for L in Γ. Then $\mathrm{C}_\Gamma(\mathrm{L}_\alpha) = \{e\}$ for all $\alpha \leqslant \lambda$. By transfinite induction each L_α is ICC. Another induction shows that each $\boldsymbol{a} \upharpoonright \mathrm{L}_\alpha$ is free: this is clear for limit α, and at successors, L_α is either normal or finite index in $\mathrm{L}_{\alpha+1}$, so assuming $\boldsymbol{a} \upharpoonright \mathrm{L}_\alpha$ is free it follows that $\boldsymbol{a} \upharpoonright \mathrm{L}_{\alpha+1}$ is free by applying either Proposition 4.11 (Proposition 4.4 also works) or Proposition 4.5.

If now L is shift-minimal and $\boldsymbol{a}$ is a non-trivial m.p. action of Γ with $\boldsymbol{a} \prec \boldsymbol{s}_\Gamma$ then $\boldsymbol{a} \upharpoonright \mathrm{L} \prec \boldsymbol{s}_\mathrm{L}$ so that $\boldsymbol{a} \upharpoonright \mathrm{L}$ is free and thus $\boldsymbol{a}$ is free.

(2): Assume that both $\boldsymbol{a} \upharpoonright \mathrm{L}$ and $\boldsymbol{a} \upharpoonright \mathrm{C}_\Gamma(\mathrm{L})$ are free. Let $\{\mathrm{L}_\alpha\}_{\alpha \leqslant \lambda}$ be an ascendant series for L in Γ. Theorem B.9 implies that $\mathrm{AR}_{\mathrm{L}_\alpha} = \mathrm{AR}_{\mathrm{C}_\Gamma(\mathrm{L}_\alpha)} = \{e\}$ for all $\alpha \leqslant \lambda$. For each $\alpha \leqslant \lambda$ we have

$$\{e\} = \mathrm{AR}_{\mathrm{C}_\Gamma(\mathrm{L}_\alpha)} \cap \mathrm{L}_{\alpha+1} = \mathrm{AR}_{\mathrm{C}_\Gamma(\mathrm{L}_\alpha)} \cap \mathrm{C}_{\mathrm{L}_{\alpha+1}}(\mathrm{L}_\alpha) = \mathrm{AR}_{\mathrm{C}_{\mathrm{L}_{\alpha+1}}(\mathrm{L}_\alpha)}$$

where the last equality follows from Corollary B.4 since the series $\{\mathrm{C}_{\mathrm{L}_\beta}(\mathrm{L}_\alpha)\}_{\beta \leqslant \lambda}$ is ascendant in $\mathrm{C}_\Gamma(\mathrm{L}_\alpha)$. It is clear that $\mathrm{C}_{\mathrm{L}_{\alpha+1}}(\mathrm{L}_\alpha) \leqslant \mathrm{C}_\Gamma(\mathrm{L})$, so by hypothesis $\boldsymbol{a} \upharpoonright \mathrm{C}_{\mathrm{L}_{\alpha+1}}(\mathrm{L}_\alpha)$ is free for all $\alpha \leqslant \lambda$. We now show by transfinite induction on $\alpha \leqslant \lambda$ that $\boldsymbol{a} \upharpoonright \mathrm{L}_\alpha$ is free. The induction is clear at limit stages. At successor stages, if we assume for induction that $\boldsymbol{a} \upharpoonright \mathrm{L}_\alpha$ is free then all the hypotheses of Proposition 4.11 hold and it follows that $\boldsymbol{a} \upharpoonright \mathrm{L}_{\alpha+1}$ is free. ☐

Proposition 4.14. *Let $\boldsymbol{a} = \Gamma \curvearrowright^{a} (X, \mu)$ be a measure preserving action of Γ. Let $\mathrm{K} = \ker(\boldsymbol{a})$.*

1. *Suppose that there exists a normal subgroup N of Γ such that $\boldsymbol{a} \upharpoonright \mathrm{N}$ is free and such that every finite index subgroups of N acts ergodically. Then $\Gamma_x = \mathrm{K}$ almost surely.*

2. *Suppose that* $\boldsymbol{a}$ *is NA-ergodic and there exists a non-amenable normal subgroup* N *of* Γ *such that* $\boldsymbol{a} \upharpoonright \mathrm{N}$ *is free. Then* K *is amenable and* $\Gamma_x = \mathrm{K}$ *almost surely.*

Proof. We begin with (1). Note that, by Proposition 4.4, if $\gamma \in \Gamma$ is such that the set $\{h\gamma h^{-1} : h \in \mathrm{N}\}$ is infinite, then $\mu(\mathrm{Fix}^a(\gamma)) = 0$. It therefore suffices to show that if $\mu(\mathrm{Fix}^a(\gamma)) > 0$ and $\{h\gamma h^{-1} : h \in \mathrm{N}\}$ is finite, then $\gamma \in \mathrm{K}$. This set being finite means that the group $\mathrm{H} = \mathrm{C}_\Gamma(\gamma) \cap \mathrm{N}$ is finite index in N, so $\boldsymbol{a} \upharpoonright \mathrm{H}$ is ergodic by hypothesis. Since $\mathrm{H} \leqslant \mathrm{C}_\Gamma(\gamma)$, the set $\mathrm{Fix}^a(\gamma)$ is $\boldsymbol{a} \upharpoonright \mathrm{H}$-invariant, so if it is non-null then it must be conull, i.e., $\gamma \in \mathrm{K}$, by ergodicity.

For (2), amenability of K is immediate since $\boldsymbol{a}$ is non-trivial and NA-ergodic. NA-ergodicity also implies that every finite index subgroup of N acts ergodically, so (1) applies and we are done. ☐

The following Corollary replaces the hypothesis in Proposition 4.13.(1) that $\mathrm{C}_\Gamma(\mathrm{L}) = \{e\}$ with the hypotheses that $\mathrm{AR}_\Gamma = \{e\}$ and $\boldsymbol{a}$ is NA-ergodic.

Corollary 4.15. *Suppose* $\mathrm{AR}_\Gamma = \{e\}$*. Let* $\boldsymbol{a}$ *be any NA-ergodic action of* Γ *and suppose that there exists a non-trivial almost ascendant subgroup* L *of* Γ *such that the restriction* $\boldsymbol{a} \upharpoonright \mathrm{L}$ *of* $\boldsymbol{a}$ *to* L *is free, then* $\boldsymbol{a}$ *itself is free.*

Proof. Let $\{\mathrm{L}_\alpha\}_{\alpha \leqslant \lambda}$ be an almost ascendant series for L in Γ. Since $\mathrm{AR}_\Gamma = \{e\}$, Corollary B.4 implies that $\mathrm{AR}_{\mathrm{L}_\alpha} = \{e\}$ for each $\alpha \leqslant \lambda$. Suppose for induction that we have shown that $\boldsymbol{a} \upharpoonright \mathrm{L}_\alpha$ is free for all $\alpha < \beta$. If β is a limit then $\mathrm{L}_\beta = \bigcup_{\alpha<\beta} \mathrm{L}_\alpha$ so $\boldsymbol{a} \upharpoonright \mathrm{L}_\beta$ is free as well. If $\beta = \alpha + 1$ is a successor then $\boldsymbol{a} \upharpoonright \mathrm{L}_\alpha$ is free and L_α is either finite index or normal in L_β. If L_α is finite index in L_β then $\boldsymbol{a} \upharpoonright \mathrm{L}_\beta$ is free by Proposition 4.5. If L_α is normal in L_β then $\boldsymbol{a} \upharpoonright \mathrm{L}_\beta$ is free by Proposition 4.14.(2). It follows by induction that $\boldsymbol{a} \upharpoonright \Gamma$ is free. ☐

Corollary 4.16.

1. *Let* Γ *be a countable group with* $\mathrm{AR}_\Gamma = \{e\}$*. If* Γ *contains a shift-minimal almost ascendant subgroup* L *then* Γ *is itself shift-minimal.*

2. *Suppose that* Γ *is a countable group containing an ascendant subgroup* L *such that* L *is shift-minimal and* $\mathrm{AR}_{\mathrm{C}_\Gamma(\mathrm{L})} = \{e\}$*. Then* Γ *is shift-minimal. In particular, if both* L *and* $\mathrm{C}_\Gamma(\mathrm{L})$ *are shift-minimal then so is* Γ*.*

Proof. Starting with (1), let L be a shift-minimal almost ascendant subgroup of Γ. Let $\boldsymbol{a}$ be a non-trivial measure preserving action of Γ weakly contained in $\boldsymbol{s}_\Gamma$. Then $\boldsymbol{a}$ is NA-ergodic and $\boldsymbol{a} \upharpoonright \mathrm{L}$ is free, so $\boldsymbol{a}$ is free by Corollary 4.15. Statement (2) is a special case of (1) since Theorem B.9 shows that $\mathrm{AR}_\Gamma = \{e\}$. ☐

5 Examples of shift-minimal groups

Theorem 5.15 below shows that if the reduced C*-algebra of a countable group Γ admits a unique tracial state then Γ is shift-minimal. We can also often gain more specific information by giving direct ergodic theoretic proofs of shift-minimality. These proofs often rely on an appeal to some form of the Poincaré recurrence theorem (several proofs of which may be found in [Ber96]).

5.1 Free groups

Since the argument is quite short it seems helpful to present a direct argument that free groups are shift-minimal.

Theorem 5.1. *Let* Γ *be a non-abelian free group.*

(i) If $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ *is any non-trivial measure preserving action of* Γ *which is NA-ergodic then* $\boldsymbol{a}$ *is free.*

(ii) Γ *is shift-minimal.*

Proof. For (i) we show that non-free actions of Γ are never NA-ergodic. Suppose that $\boldsymbol{a}$ is non-free so that $\mu(\mathrm{Fix}^a(\gamma)) > 0$ for some $\gamma \in \Gamma - \{e\}$. Fix any $\delta \in \Gamma - \langle\gamma\rangle$. By the Poincaré recurrence theorem there exists an $n > 0$ with $\mu(\delta^n \cdot \mathrm{Fix}^a(\gamma) \cap \mathrm{Fix}^a(\gamma)) > 0$. The group H generated by $\delta^n\gamma\delta^{-n}$ and γ is free on these elements and $\alpha^a \cdot x = x$ for every $\alpha \in H$ and $x \in \delta^n \cdot \mathrm{Fix}^a(\gamma) \cap \mathrm{Fix}^a(\gamma)$. In particular $\boldsymbol{a} \upharpoonright H$ is not ergodic, whence $\boldsymbol{a}$ cannot be NA-ergodic.

Statement (ii) now follows since any non-trivial action weakly contained in $\boldsymbol{s}_\Gamma$ is strongly NA-ergodic, hence free by (i). ☐

Another proof of part (i) of Theorem 5.1 follows from Theorem 3.13 (see also [AGV12, Lemma 24]). Indeed, alternative (2) of Theorem 3.13 can never hold since a non-abelian free group has only countably many amenable subgroups. So if $\boldsymbol{a}$ is any non-trivial NA-ergodic action of a non-abelian free group Γ then (1) of Theorem 3.13 holds, and so $\boldsymbol{a}$ is free since the only normal amenable subgroup of Γ is the trivial group $N = \{e\}$.

5.2 Property (BP)

Definition 5.2. Let Γ be a countable group.

1. Γ is said to be a *Powers group* ([Ha85]) if $\Gamma \neq \{e\}$ and for every finite subset $F \subseteq \Gamma \setminus \{e\}$ and every integer $N > 0$ there exists a partition $\Gamma = C \sqcup D$ and elements $\alpha_1, \dots, \alpha_N \in \Gamma$ such that

$$\gamma C \cap C = \varnothing \quad \text{for all } \gamma \in F$$
$$\alpha_j D \cap \alpha_k D = \varnothing \quad \text{for all } j, k \in \{1, \dots, N\},\ j \neq k.$$

   Γ is said to be a *weak Powers group* ([BN88]) if Γ satisfies all instances of the Powers property with F ranging over finite subsets of mutually conjugate elements of $\Gamma\setminus\{e\}$. We define Γ to be a *weak* Powers group* if Γ satisfies all instances of the Powers property with F ranging over singletons in $\Gamma \setminus \{e\}$.

2. Γ has *property* P_{nai} ([BCH94]) if for any finite subset F of Γ there exists an element $\alpha \in \Gamma$ of infinite order such that for each $\gamma \in F$, the canonical homomorphism from the free product $\langle\gamma\rangle * \langle\alpha\rangle$ onto the subgroup $\langle\gamma, \alpha\rangle$ of Γ generated by γ and α is an isomorphism.

   If Γ satisfies the defining property of P_{nai} but with F only ranging over singletons, then we say that Γ has *property* P^*_{nai}.

3. Γ is said to have property (PH) ([Pro93]) if for all nonempty finite $F \subseteq \Gamma\setminus\{e\}$ there exists some ordering $F = \{\gamma_1, \dots, \gamma_m\}$ of F along with an increasing sequence $e \in Q_1 \subseteq \cdots \subseteq Q_m$ of subsets of Γ such that for all $i \leqslant m$, all nonempty finite $M \subseteq Q_i$ and all $n > 0$ we may find $\alpha_1, \dots, \alpha_n \in Q_i$ and $T_1, \dots, T_n$ pairwise disjoint such that

$$(\alpha_j\delta)\gamma_i(\alpha_j\delta)^{-1}(\Gamma \setminus T_j) \subseteq T_j$$

   for all $\delta \in M$ and $1 \leqslant j \leqslant n$.

Examples of groups with these properties may be found in [AM07, HP11, MOY11, PT11] along with the references given in the above definitions. For our purposes, what is important is a common consequence of these properties.

Definition 5.3. A countable group Γ is said to have property (BP) if for all $\gamma \in \Gamma \setminus \{e\}$ and $n \geqslant 2$ there exists $\alpha_1, \dots, \alpha_n \in \Gamma$, a subgroup $H \leqslant \Gamma$, and pairwise disjoint subsets $T_1, \dots, T_n \subseteq H$ such that

$$\alpha_j \gamma \alpha_j^{-1}(H \setminus T_j) \subseteq T_j$$

for all $j = 1, \dots, n$.

Note that when γ, H, $\alpha_1, \dots, \alpha_n$, and $T_1, \dots, T_n$ are as above, then $\alpha_j \gamma \alpha_j^{-1} \in H$ and $T_j \neq \varnothing$ for all $j \leqslant n$.

We show in Theorem 5.6 that groups with property (BP) satisfy a strong form of shift-minimality. The definition of property (BP) (as well as its name) is motivated by an argument of M. Brin and G. Picioroaga showing that all weak Powers groups contain a free group. Their proof appears in [Ha07] (see the remark following Question 15 in that paper), though we also present a version of their proof in Theorem 5.4 since we will need it for Theorem 5.6.

Theorem 5.4 (Brin, Picioroaga [Ha07])**.**

(1) All weak* Powers groups have property (BP)*.*

(2) Property P^*_{nai} implies property (BP)*.*

(3) Property (PH) *implies property* (BP)*.*

(4) Groups with property (BP) *contain a free group.*

Proof. (1): given $\gamma \in \Gamma \setminus \{e\}$ and $n \geqslant 1$ by the weak* Powers property there exists $\alpha_1, \dots, \alpha_n$ and a partition $\Gamma = C \sqcup D$ of Γ with $\gamma C \cap C = \varnothing$ and $\alpha_i D \cap \alpha_j D = \varnothing$ for all $1 \leqslant i, j \leqslant n$, $i \neq j$. Take $H = \Gamma$ and for each $1 \leqslant j \leqslant n$ let $T_j = \alpha_j D$ so that the sets $T_1, \dots, T_n$ are pairwise disjoint and

$$\alpha_j \gamma \alpha_j^{-1}(\Gamma \setminus T_j) = \alpha_j \gamma (\Gamma \setminus D) = \alpha_j \gamma C \subseteq \alpha_j(\Gamma \setminus C) = \alpha_j D = T_j$$

thus verifying (BP).

(2): Let $\gamma \in \Gamma \setminus \{e\}$. By property P^*_{nai} there exists an element $\alpha \in \Gamma$ of infinite order such that the subgroup $H = \langle \gamma, \alpha \rangle$ of Γ is canonically isomorphic to the free product $\langle \gamma \rangle * \langle \alpha \rangle$. Let T_n denote the set of elements of H whose reduced expression starts with $\alpha^n \gamma^k$ for some $k \in \mathbb{Z}$ with $\gamma^k \neq e$. Then the sets T_n, $n \in \mathbb{N}$, are pairwise disjoint and $\alpha^n \gamma \alpha^{-n}(H \setminus T_n) \subseteq T_n$.

(3): Assume that Γ has property (PH) and fix any $\gamma \in \Gamma \setminus \{e\}$ and $n \geqslant 1$ toward the aim of verifying property (BP). Taking $F = \{\gamma\}$ we obtain a set $Q = Q_1 \subseteq \Gamma$ from the above definition of (PH) with $e \in Q$. Taking $M = \{e\}$, the defining property of Q produces $\alpha_1, \dots, \alpha_n \in Q$ and pairwise disjoint $T_1, \dots, T_n \subseteq \Gamma$ with

$$\alpha_j \gamma \alpha_j^{-1}(\Gamma \setminus T_j) \subseteq T_j,$$

so taking $H = \Gamma$ confirms this instance of property (BP).

Statement (4) is a consequence of the following Lemma, which will be used in Theorem 5.6 below.

Lemma 5.5 (Brin, Picioroaga)**.** *Suppose that $x_1, \dots x_4$ are elements of a group H and that $T_1, \dots, T_4$ are pairwise disjoint subsets of H such that*

$$x_j(H \setminus T_j) \subseteq T_j$$

for each $j \in \{1, \dots, 4\}$. Then the group elements $u = x_1 x_2$ and $v = x_3 x_4$ freely generate a non-abelian free subgroup of H.

Proof of Lemma 5.5. The hypothesis $x_j(H \setminus T_j) \subseteq T_j$ implies that also $x_j^{-1}(H \setminus T_j) \subseteq T_j$. For distinct $i, j \in \{1, \dots, 4\}$ it then follows that

$$\begin{aligned} x_i x_j(H \setminus T_j) &\subseteq x_i T_j \subseteq x_i(H \setminus T_i) \subseteq T_i \\ \text{and} \quad (x_i x_j)^{-1}(H \setminus T_i) &\subseteq x_j^{-1} T_i \subseteq x_j^{-1}(H \setminus T_j) \subseteq T_j \end{aligned}$$

so for $u = x_1 x_2$ and $v = x_3 x_4$ we have

$$\begin{aligned} u(H \setminus T_2) &\subseteq T_1 \quad & u^{-1}(H \setminus T_1) &\subseteq T_2 \\ v(H \setminus T_4) &\subseteq T_3 \quad & v^{-1}(H \setminus T_3) &\subseteq T_4. \end{aligned}$$

A ping pong argument now shows that u and v freely generate a non-abelian free subgroup of H. ☐[Lemma 5.5]

If now Γ has property (BP) then taking any $\gamma \in \Gamma \setminus \{e\}$ and $n = 4$ we obtain $\alpha_1, \dots, \alpha_4 \in \Gamma$, $H \leqslant \Gamma$ and $T_1, \dots, T_4 \subseteq H$ as in the definition of property (BP). Lemma 5.5 now applies with $x_j = \alpha_j \gamma \alpha_j^{-1}$ for $j \in \{1, \dots, 4\}$. ☐[Theorem 5.4]

Lemma 5.5 can be used to show that any non-trivial ergodic invariant random subgroup of a group with property (BP) contains a free group.

Theorem 5.6. *Let Γ have property* (BP) *and let $\boldsymbol{a} = \Gamma \curvearrowright^a (Y, \nu)$ be an ergodic measure preserving action of Γ. Suppose that $\boldsymbol{a}$ is non-free. Then the stabilizer of ν-almost every $y \in Y$ contains a non-abelian free group. In particular, all groups with property* (BP) *are shift-minimal.*

Proof. Since $\boldsymbol{a}$ is non-free there exists an element $\gamma \in \Gamma \setminus \{e\}$ such that $\nu(A) = r > 0$ where $A = \mathrm{Fix}^a(\gamma)$. By the Poincaré recurrence theorem, for all large enough n (depending on r), if $A_1, \dots, A_n \subseteq Y$ is any sequences of measurable subsets of Y each of measure r, then there exist distinct $i_1, \dots, i_4 \leqslant n$ with $\nu(A_{i_1} \cap A_{i_2} \cap A_{i_3} \cap A_{i_4}) > 0$. Pick such an n with $n \geqslant 4$. By property (BP) there exists $\alpha_1, \dots, \alpha_n \in \Gamma$, $H \leqslant \Gamma$, and pairwise disjoint $T_1, \dots, T_n \subseteq H$ such that $\alpha_i \gamma \alpha_i^{-1}(H \setminus T_i) \subseteq T_i$ for all $i \in \{1, \dots, n\}$. By our choice of n there must exist distinct $i_1, \dots, i_4 \leqslant n$ such that

$$\nu(\alpha_{i_1}^a A \cap \alpha_{i_2}^a A \cap \alpha_{i_3}^a A \cap \alpha_{i_4}^a A) > 0. \tag{5.1}$$

For $j = 1, \dots, 4$ let $x_j = \alpha_{i_j} \gamma \alpha_{i_j}^{-1}$. Lemma 5.5 (applied to $x_1, \dots x_4$ and $T_1, \dots T_4$) shows that $\langle x_1, \dots, x_4 \rangle$ contains a free group. Additionally, (5.1) shows that $\nu(\mathrm{Fix}^a(\langle x_1, \dots, x_4 \rangle)) > 0$ since

$$\mathrm{Fix}^a(\langle x_1, \dots, x_4 \rangle) \supseteq \bigcap_{j=1}^{4} \mathrm{Fix}^a(x_i) = \bigcap_{j=1}^{4} \alpha_{i_j}^a A.$$

The event that Γ_y contains a free group is therefore non-null. This event is also $\boldsymbol{a}$-invariant, so ergodicity now implies that almost every stabilizer contains a free group.

If now $\boldsymbol{b}$ is any non-trivial measure preserving action of Γ weakly contained in $\boldsymbol{s}_\Gamma$ then $\boldsymbol{b}$ is ergodic and by Lemma 3.11 almost every stabilizer is amenable hence does not contain a free group. Then $\boldsymbol{b}$ is essentially free by what we have already shown. Therefore Γ is shift-minimal. ☐

In [Be91] Bèdos defines a group Γ to be an *ultraweak Powers group* if it has a normal subgroup N that is a weak Powers group such that $C_\Gamma(N) = \{e\}$. Let us say that Γ is an *ultraweak* Powers group* if it has a normal subgroup N that is an weak* Powers group such that $C_\Gamma(N) = \{e\}$.

Theorem 5.7. *Let Γ be a countable group.*

1. *Suppose that* Γ *contains an almost ascendant subgroup* L *with property* (BP) *such that* $C_\Gamma(L) = \{e\}$*. Then for every ergodic m.p. action* $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ *of* Γ*, either* $\boldsymbol{a}$ *is free or* $\Gamma_x \cap L$ *contains a non-abelian free group almost surely.*
2. *Suppose that* Γ *contains an ascendant subgroup* L *such that both* L *and* $C_\Gamma(L)$ *have property* (BP)*. Then for every ergodic m.p. action* $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ *of* Γ*, either* $\boldsymbol{a}$ *is free,* $\Gamma_x \cap L$ *contains a non-abelian free group almost surely, or* $\Gamma_x \cap C_\Gamma(L)$ *contains a non-abelian free group almost surely.*
3. *Every non-trivial ergodic invariant random subgroup of an ultraweak***-Powers group contains a non-abelian free group almost surely.*

Proof. (1) Since L has property (BP) it is ICC, so if $\boldsymbol{a} \upharpoonright L$ is free then $\boldsymbol{a}$ itself is free by part (1) of Proposition 4.13. Suppose then that $\boldsymbol{a} \upharpoonright L$ is non-free. Let $\pi : (X, \mu) \to (Z, \eta)$ be the ergodic decomposition map for $\boldsymbol{a} \upharpoonright L$ and let $\mu = \int_z \mu_z \, d\eta(z)$ be the disintegration of μ with respect to η. Since $\boldsymbol{a} \upharpoonright L$ is non-free then the set $A \subseteq Z$, consisting of of all $z \in Z$ such that $L \curvearrowright^a (X, \mu_z)$ is non-free, is η-non-null. If $z \in A$ then $\mu_z(\{x : L_x \text{ contains a non-abelian free group}\}) = 1$ by Theorem 5.6. The event that L_x contains a non-abelian free group is therefore μ-non-null. This event is Γ-invariant (a subgroup contains a free group if and only if any of its conjugates contains one), so ergodicity implies that L_x contains a free group almost surely. Since $L_x = \Gamma_x \cap L$ we are done.

The proof of (2) is similar, using part (2) of Proposition 4.13. (3) is immediate from (1) and the definitions. ☐

We note also that (BP) is preserved by extensions.

Proposition 5.8. *Let* N *be a normal subgroup of* Γ*. If* N *and* Γ/N *both have property* (BP) *then* Γ *also has property* (BP)*.*

Proof. Let $\gamma \in \Gamma \setminus \{e\}$ and $n \geqslant 1$ be given.

If $\gamma \in N$ then property (BP) for N implies that there exists $\alpha_1, \dots, \alpha_n \in N$, $H \leqslant N$ and pairwise disjoint $T_1, \dots, T_n \subseteq H$ as in the definition of (BP) for N. These also satisfy this instance of property (BP) for Γ.

If $\gamma \notin N$ then the image of γ in Γ/N is not the identity element so property (BP) for Γ/N implies that there exist cosets $\alpha_1 N, \cdots \alpha_n N \in \Gamma/N$, a subgroup $K \leqslant \Gamma$ containing N, and pairwise disjoint $T_1, \dots, T_n \subseteq K/N$ as in the definition of (BP) for Γ/N. Then $\alpha_1, \dots, \alpha_n$, K, and the sets $T_i' = \bigcup T_i$, $i = 1, \dots, M$, verify this instance of property (BP) for Γ. ☐

Remark 5.9. If a group Γ has property (BP) then it has the unique trace property. A quick proof of this follows [BCH94]. The proof of this is almost exactly as in [BCH94, Lemma 2.2] with just a minor adjustment to the first part of their proof which we now describe. One first shows for any $\gamma \in \Gamma \setminus \{e\}$ and any $n \geqslant 2$, if $\alpha_1, \dots, \alpha_n$, H, and $T_1, \dots, T_n$ are as in the definition of (BP) then for all $z = (z_1, \dots, z_n) \in \mathbb{C}^n$ we have

$$\Big|\Big| \sum_{j=1}^{n} z_j \lambda_\Gamma(\alpha_j \gamma \alpha_j^{-1}) \Big|\Big| \leqslant 2 ||z||_2. \tag{5.2}$$

Let $x_j = \alpha_j \gamma \alpha_j^{-1}$ so that $x_j \in H$ and $x_j(H \setminus T_j) \subseteq T_j$ for all $j = 1, \dots, n$. Let 1_A denote the indicator function of a subset $A \subseteq H$. For $f, g \in \ell^2(H)$ we then have

$$\begin{aligned} |\langle \lambda_H(x_j) f, g\rangle| &\leqslant |\langle \lambda_H(x_j)(1_{T_j} f), g\rangle| + |\langle \lambda_H(x_j)(1_{H \setminus T_j} f), g\rangle| \\ &= |\langle \lambda_H(x_j)(1_{T_j} f), g\rangle| + |\langle 1_{x_j(H \setminus T_j)} \lambda_H(x_j)(f), 1_{T_j} g\rangle| \leqslant ||1_{T_j} f|| \, ||g|| + ||f|| \, ||1_{T_j} g||. \end{aligned}$$

The remainder of the proof of (5.2) now proceeds as in [BCH94, Lemma 2.2] using that the T_j are pairwise disjoint. It now follows as in the paragraph following [BCH94, Definition 1] that $C_r^*(\Gamma)$ has a unique tracial state.

5.3 Linear groups

In the case that Γ is a countable linear group, a theorem of Y. Glasner [Gl12] shows that the existence of a non-trivial normal amenable subgroup is the only obstruction to shift-minimality: Glasner shows that every amenable invariant random subgroup of a linear group Γ must concentrate on the subgroups of the amenable radical of Γ. Along with Proposition 3.15 this implies that a countable linear group Γ is shift-minimal if and only if Γ contains no non-trivial normal amenable subgroups. Another way to deduce these results is to use Theorem 5.14 below along with the following Theorem of Poznansky.

Theorem 5.10 (Theorem 1.1 of [Poz09]). *Let Γ be a countable linear group. Then the following are equivalent*

(1) Γ is C^*-simple.

(2) Γ has the unique trace property.

(3) Γ contains no non-trivial normal amenable subgroups, i.e., $\mathrm{AR}_\Gamma = \{e\}$.

Corollary 5.11. *Let Γ be a countable linear group. The properties (1), (2), and (3) of Theorem 5.10 are equivalent to each of the following properties:*

(4) Γ is shift-minimal.

(5) Γ has no non-trivial amenable invariant random subgroups.

Proof. The implication (2)⇒(5) follows from Theorem 5.14, the implication (5)⇒(4) is Corollary 3.14, and (4)⇒(3) follows from Proposition 3.15. The remaining implications follow from Poznansky's Theorem 5.10. ☐

5.4 Unique tracial state on $C^*_r(\Gamma)$

We write $C^*_r(\Gamma)$ for the reduced C^*-algebra of Γ. This is the C^*-algebra generated by $\{\lambda_\Gamma(\gamma) : \gamma \in \Gamma\}$ in $\mathcal{B}(\ell^2(\Gamma))$, where λ_Γ denotes the left regular representation of Γ. Let $1_e \in \ell^2(\Gamma)$ denote the indicator function of $\{e\}$. We obtain a tracial state τ_Γ, called the *canonical trace* on $C^*_r(\Gamma)$, given by $\tau_\Gamma(a) = \langle a(1_e), 1_e\rangle$.

Let ρ be a probability measure on Sub_Γ and define the function $\varphi_\rho \in \ell^\infty(\Gamma)$ by

$$\varphi_\rho(\gamma) = \rho(\{H : \gamma \in H\}).$$

It is shown in [IKT09] (see also Theorem A.16) and [Ve11] that φ_ρ is a positive definite function on Γ. It will be useful here to identify φ_ρ as the diagonal matrix coefficient of a specific unitary representation of Γ described below.

Consider the field of Hilbert spaces $\{\ell^2(\Gamma/H) : H \in \mathrm{Sub}_\Gamma\}$. For $\gamma \in \Gamma$ denote by $x^\gamma \in \prod_H \ell^2(\Gamma/H)$ the vector field $x^\gamma_H = 1_{\gamma H}$ where $1_{\gamma H} \in \ell^2(\Gamma/H)$ is the indicator function of the singleton set $\{\gamma H\} \subseteq \Gamma/H$. Then $\{x^\gamma\}_{\gamma\in\Gamma}$ determines a fundamental family of measurable vector fields and we let $\mathcal{H}_\rho = \int_H^\oplus \ell^2(\Gamma/H)\,d\rho$ denote the corresponding Hilbert space consisting of all square integrable measurable vector fields. The inner product on $\mathcal{H}_\rho$ is given by $\langle x, y\rangle = \int_H \langle x_H, y_H\rangle_{\ell^2(\Gamma/H)}\,d\rho$. Define the unitary representation λ_ρ of Γ on $\mathcal{H}_\rho$ by

$$\lambda_\rho = \int_H^\oplus \lambda_{\Gamma/H}\,d\rho,$$

i.e., $\lambda_\rho(\gamma)(x)_H = \lambda_{\Gamma/H}(\gamma)(x_H)$, where $\lambda_{\Gamma/H}$ denotes the quasi-regular representation of Γ on $\ell^2(\Gamma/H)$. We then have

$$\begin{aligned}\langle\lambda_\rho(\gamma)(x^e), x^e\rangle &= \int_H \langle\lambda_\rho(\gamma)(x^e)_H, x^e_H\rangle_{\ell^2(\Gamma/H)}\, d\rho \\ &= \int_H \langle\lambda_{\Gamma/H}(\gamma)(1_H), 1_H\rangle_{\ell^2(\Gamma/H)}\, d\rho = \rho(\{H : \gamma\in H\}) = \varphi_\rho(\gamma).\end{aligned}$$

We have shown the following.

Proposition 5.12. *$(\mathcal{H}_\rho, \lambda_\rho, x^e)$ is the* GNS *triple associated with the positive definite function φ_ρ on Γ.*

It is clear that if ρ is conjugation invariant (i.e., if ρ is an invariant random subgroup) then φ_ρ will be constant on each conjugacy class of Γ.

Lemma 5.13. *If H is an amenable subgroup of Γ then $\lambda_{\Gamma/H}$ is weakly contained in λ_Γ. Thus, for all $f\in\ell^1(\Gamma)$ we have $\|\lambda_{\Gamma/H}(f)\| \leqslant \|\lambda_\Gamma(f)\|$.*

Proof. H being amenable implies that the trivial one dimensional representation 1_H of H is weakly contained in the left regular representation λ_H of H ([BHV08, Theorem G.3.2]). Thus by [BHV08, Theorem F.3.5] we have $\lambda_{\Gamma/H} \cong \mathrm{Ind}_H^\Gamma(1_H) \prec \mathrm{Ind}_H^\Gamma(\lambda_H) \cong \lambda_\Gamma$. The second statement follows immediately from [BHV08, F.4.4]. ☐

Theorem 5.14. *If ρ is any measure on Sub_Γ concentrating on the amenable subgroups then λ_ρ is weakly contained in the left regular representation λ_Γ of Γ.*

Therefore, if θ is an amenable invariant random subgroup of Γ then φ_θ extends to a tracial state on $C^*_r(\Gamma)$ which is distinct from the canonical trace τ_Γ whenever θ is non-trivial.

Proof. By [BHV08, F.4.4] to show that $\lambda_\rho \prec \lambda_\Gamma$ it suffices to show that $\|\lambda_\rho(f)\| \leqslant \|\lambda_\Gamma(f)\|$ for all $f\in\ell^1(\Gamma)$. Using that ρ concentrates on the amenable subgroups and Lemma 5.13 we have for $f\in\ell^1(\Gamma)$ and $x, y\in\mathcal{H}_\rho$

$$\begin{aligned}|\langle\lambda_\rho(f)x, y\rangle| &= \Big|\int_H \langle\lambda_{\Gamma/H}(f)(x_H), y_H\rangle_{\ell^2(\Gamma/H)}\, d\rho\Big| \\ &\leqslant \int_H \|\lambda_{\Gamma/H}(f)\|\, \|x_H\|\, \|y_H\|\, d\rho \\ &\leqslant \|\lambda_\Gamma(f)\| \int_H \|x_H\|\, \|y_H\|\, d\rho \\ &\leqslant \|\lambda_\Gamma(f)\|\, \|x\|\, \|y\|\end{aligned}$$

from which we conclude that $\|\lambda_\rho(f)\| \leqslant \|\lambda_\Gamma(f)\|$.

Suppose now θ is an amenable invariant random subgroup of Γ. Since λ_θ is weakly contained in λ_Γ, λ_θ extends to a representation of $C^*_r(\Gamma)$ and φ_θ extends to a state on $C^*_r(\Gamma)$ via $a \mapsto \langle\lambda_\theta(a)(x^e), x^e\rangle$. Since φ_θ is conjugation invariant this is a tracial state. If θ is non-trivial then there is some $\gamma\in\Gamma\setminus\{e\}$ with $\varphi_\theta(\gamma) = \theta(\{H : \gamma\in H\}) > 0$ showing that this is distinct from the canonical trace. ☐

Corollary 5.15. *Let Γ be a countable group with the unique trace property. Then Γ has no non-trivial amenable invariant random subgroups. It follows that every non-trivial NA-ergodic action of Γ is free and Γ is shift-minimal.*

Proof. That Γ has no non-trivial amenable invariant random subgroups follow from Theorem 5.14. If $\boldsymbol{a}$ is a non-trivial NA-ergodic action of Γ then the invariant random subgroup $\theta_{\boldsymbol{a}}$ is amenable by Theorem 3.13, and thus $\theta_{\boldsymbol{a}} = \delta_e$, i.e., $\boldsymbol{a}$ is free. Since every m.p. action weakly contained in $\boldsymbol{s}_\Gamma$ is NA-ergodic, Γ is also shift-minimal. ☐

Remark 5.16. The positive definite function φ_θ associated to an invariant random subgroup θ is also realized in the Koopman representation $\kappa_0^{\boldsymbol{s}_\theta}$ corresponding to the θ-random Bernoulli shift $\boldsymbol{s}_{\theta,\eta}$ of Γ with a non-atomic base space (Z, η) (see [T-D12a] for the definition of the θ-random Bernoulli shift). Indeed, take $Z = \mathbb{R}$ and take η to be the standard Gaussian measure (with unit variance). Let $p_\gamma : \mathbb{R}^{\leqslant\backslash\Gamma} \to \mathbb{R}$ be the function $p_\gamma(f) = f(H_f\gamma)$. Then $p_\gamma \in L^2_0(\eta^{\theta\backslash\Gamma})$ and each p_γ is a unit vector. In addition we have $\kappa_0^{\boldsymbol{s}_{\theta,\eta}}(\gamma)(p_e) = p_\gamma$ and

$$\langle p_\gamma, p_e\rangle = \int_H \int_{f\in\mathbb{R}^{H\backslash\Gamma}} f(H\gamma)f(H)\, d\eta^{H\backslash\Gamma}\, d\theta(H) = \int_H 1_{\{H\,:\,H\gamma=H\}}\, d\theta = \varphi_\theta(\gamma) \tag{5.3}$$

and so $(L^2_0(\eta^{\theta\backslash\Gamma}), \kappa_0^{\boldsymbol{s}_{\theta,\eta}}, p_e)$ is a triple realizing φ_θ.

6 Cost

6.1 Notation and background

See [Ga00] and [KM05] for background on the theory of cost of equivalence relations and groups. We recall the basic definitions to establish notation and terminology.

Definition 6.1. Let (X, μ) be a standard non-atomic probability space.

(i) By an *L-graphing* on (X, μ) we mean a countable collection $\Phi = \{\varphi_i : A_i \to B_i\}_{i\in I}$ of partial Borel automorphism of X that preserve the measure μ. The *cost* of the L-graphing Φ is given by

$$C_\mu(\Phi) = \sum_{i\in I} \mu(A_i).$$

In (ii)-(vi) below Φ denotes an L-graphing on (X, μ).

(ii) We denote by $\mathcal{G}_\Phi$ the graph on X associated to Φ, i.e., for $x, y \in X$, $(x, y) \in \mathcal{G}_\Phi$ if and only if $x \neq y$ and $\varphi^{\pm1}(x) = y$ for some $\varphi \in \Phi$. We let $d_\Phi : X \times X \to \mathbb{N} \cup \{\infty\}$ denote the graph distance corresponding to $\mathcal{G}_\Phi$, i.e., for $x, y \in X$,

$$d_\Phi(x, y) = \inf\{m \in \mathbb{N} : \exists\varphi_0, \dots, \varphi_{m-1} \in \Phi^* \ (\varphi_{m-1}^{\pm1} \circ \dots \circ \varphi_1^{\pm1} \circ \varphi_0^{\pm1}(x) = y)\}$$

where $\Phi^* = \Phi \cup \{\mathrm{id}_X\}$ and $\mathrm{id}_X : X \to X$ is the identity map.

(iii) We let E_Φ denote the equivalence relation on X generated by Φ, i.e., $xE_\Phi y \Leftrightarrow d_\Phi(x, y) < \infty$. Then E_Φ is a countable Borel equivalence relation that preserves the measure μ.

(iv) Let E be a measure preserving countable Borel equivalence relation on (X, μ). We say that Φ is an *L-graphing of* E if there is a conull set $X_0 \subseteq X$ such that $E_\Phi \upharpoonright X_0 = E \upharpoonright X_0$. This is equivalent to the condition that $[x]_{E_\Phi} = [x]_E$ for μ-almost every $x \in X$. The *cost* of E is defined as

$$C_\mu(E) = \inf\{C_\mu(\Psi) : \Psi \text{ is an L-graphing of } E\}.$$

(v) Let $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ be a measure preserving action of Γ. Let Q be a subset of Γ and let $A : Q \to \mathrm{MALG}_\mu$ be a function assigning to each $\delta \in Q$ a measurable subset A_δ of X. Then a and A define an L-graphing $\Phi^{a,A} = \{\varphi_\delta^{a,A} : \delta \in Q\}$, where $\varphi_\delta^{a,A} = \delta^a \upharpoonright A_\delta$, i.e., $\mathrm{dom}(\varphi_\delta^{a,A}) = A_\delta$ and $\varphi_\delta^{a,A}(x) = \delta^a x$ for each $x \in A_\delta$. It is clear that $E_{\Phi^{a,A}} \subseteq E_a$ and

$$C_\mu(\Phi^{a,A}) = \sum_{\delta\in Q} \mu(A_\delta)$$

so that $C_\mu(\Phi^{a,A})$ only depends on the assignment A and not on the action a.

(vi) As a converse to (v), whenever $E_\Phi \subseteq E_a$ we may find a function $A = A^{a,\Phi} : \Gamma \to \mathrm{MALG}_\mu$ such that $\mathcal{G}_{\Phi^{a,A}} = \mathcal{G}_\Phi$ and $C_\mu(\Phi^{a,A}) \leqslant C_\mu(\Phi)$. Indeed, for each $\varphi \in \Phi$ there exists a measurable partition $X = \bigsqcup_{\delta\in\Gamma} A^{a,\varphi}_\delta$ such that $\varphi \upharpoonright A^{a,\varphi}_\delta = \delta^a \upharpoonright A^{a,\varphi}_\delta$. Then taking $A_\delta = \bigcup_{\varphi\in\Phi} A^{a,\varphi}_\delta$ works.

For a measure preserving action $\boldsymbol{a} = \Gamma \curvearrowright^a (X,\mu)$ of Γ denote by E_a the orbit equivalence relation generated by a. The cost of $\boldsymbol{a}$ is defined by $C(\boldsymbol{a}) = C_\mu(E_a)$. Denote by $C(\Gamma)$ the cost of the group Γ, i.e., $C(\Gamma)$ is the infinimum of costs of free m.p. actions of Γ.

By "subequivalence relation" we will always mean "Borel subequivalence relation."

6.2 Cost and weak containment in infinitely generated groups

Lemma 6.2 together with Theorem 6.4 provide a generalization of [Ke10, Theorem 10.13]. The purpose of Lemma 6.2 is to isolate versions of a few key observations from Kechris's proof.

Lemma 6.2. *Let $F \subseteq \Gamma$ be finite and let $r \in \mathbb{R}\cup\{\infty\}$. Then the following are equivalent for a measure preserving action $\boldsymbol{a} = \Gamma \curvearrowright^a (X,\mu)$ of Γ:*

1. *There exists a sub-equivalence relation E of E_a such that $E_{a\upharpoonright\langle F\rangle} \subseteq E \subseteq E_a$ and $C_\mu(E) < r$.*
2. *There exists a finite $Q \subseteq \Gamma$ containing F and a sub-equivalence relation E of E_a such that $E_{a\upharpoonright\langle F\rangle} \subseteq E \subseteq E_{a\upharpoonright\langle Q\rangle}$ and $C_\mu(E) < r$.*
3. *There exists a finite $Q \subseteq \Gamma$ containing F, an assignment $A : Q \to \mathrm{MALG}_\mu$, and a natural number $M \in \mathbb{N}$ such that*
$$C_\mu(\Phi^{a,A}) + \sum_{\gamma\in F} \mu(\{x : d_{\Phi^{a,A}}(x,\gamma^a x) > M\}) < r.$$

Proof of Lemma 6.2. We begin with the implication (3)⇒(2). If such an $A : Q \to \mathrm{MALG}_\mu$ and $M \in \mathbb{N}$ exist then define $B : Q \to \Gamma$ by taking $B \upharpoonright Q\setminus F = A \upharpoonright Q\setminus F$ and for $\gamma \in F$ taking
$$B_\gamma = A_\gamma \cup \{x : d_{\Phi^{a,A}}(x,\gamma^a x) > M\}.$$

Let $E = E_{\Phi^{a,B}}$. Then $C_\mu(E) \leqslant C_\mu(\Phi^{a,B}) < r$ and $E_{\Phi^{a,B}} \subseteq E_{a\upharpoonright\langle Q\rangle}$. In addition we have $E_{a\upharpoonright\langle F\rangle} \subseteq E_{\Phi^{a,B}}$ since for each $\gamma \in F$ and $x \in X$, either $d_{\Phi^{a,A\upharpoonright Q}}(x,\gamma^a x) \leqslant M$ so that $(x,\gamma^a x) \in E_{\Phi^{a,A}} \subseteq E_{\Phi^{a,B}}$, or $d_{\Phi^{a,A\upharpoonright Q}}(x,\gamma^a x) > M$, in which case $x \in \mathrm{dom}(\varphi^{a,B}_\gamma)$ and so $(x,\gamma^a x) \in E_{\Phi^{a,B}}$.

(2)⇒(1) is obvious, and it remains to show (1)⇒(3). Let E be as in (1) and let Φ be an L-graphing of E with $C_\mu(\Phi) = s < r$. Since $E \subseteq E_a$ we may by 6.1.(vi) assume without loss of generality that $\Phi = \Phi^{a,B}$ for some $B : \Gamma \to \mathrm{MALG}_\mu$, $\gamma \mapsto B_\gamma$. Let $\epsilon > 0$ be such that $s + \epsilon < r$.

We have $E_{a\upharpoonright\langle F\rangle} \subseteq E = E_{\Phi^{a,B}}$ so, as F is finite, if we take a large enough finite set $Q \subseteq \Gamma$ containing F, we can ensure that
$$\sum_{\gamma\in F} \mu(\{x : d_{\Phi^{a,B\upharpoonright Q}}(x,\gamma^a x) = \infty\}) < \epsilon.$$

So if we take $M \in \mathbb{N}$ large enough then
$$\sum_{\gamma\in F} \mu(\{x : d_{\Phi^{a,B\upharpoonright Q}}(x,\gamma^a x) > M\}) < \epsilon.$$

It follows that $A = B \upharpoonright Q$ and M satisfy the desired properties. □[Lemma 6.2]

Definition 6.3. For each finite $F \subseteq \Gamma$ and $r \in \mathbb{R}\cup\{\infty\}$ let $A_{F,r} = A_{F,r}(\Gamma, X, \mu)$ denote the set of $\boldsymbol{a} \in A(\Gamma, X, \mu)$ that satisfy any – and therefore all – of the equivalent properties (1)-(3) of Lemma 6.2.

It is clear that the set $A_{F,r}(\Gamma, X, \mu)$ is an isomorphism-invariant (and in fact, orbit-equivalence-invariant) subset of $A(\Gamma, X, \mu)$. In what follows, we let $\mathrm{FR}(\Gamma, X, \mu)$ denote the subset of $A(\Gamma, X, \mu)$ consisting of all free actions.

Theorem 6.4. *Let Γ be an infinite countable group. For each finite $F \subseteq \Gamma$ and $r \in \mathbb{R} \cup \{\infty\}$ the set $A_{F,r}(\Gamma, X, \mu) \cap \mathrm{FR}(\Gamma, X, \mu)$ is contained in the interior of $A_{F,r}(\Gamma, X, \mu)$. In particular, $A_{F,r}(\Gamma, X, \mu) \cap \mathrm{FR}(\Gamma, X, \mu)$ is open in $\mathrm{FR}(\Gamma, X, \mu)$.*

Proof. Let $\boldsymbol{a} \in A_{F,r}$ be free and let $Q \subseteq \Gamma$, $A : Q \to \mathrm{MALG}_\mu$ and $M \in \mathbb{N}$ be given by Lemma 6.2.(3). For each $\gamma \in F$ let $s^a_\gamma = \mu(\{x : d_{\Phi^{a,A}}(x, \gamma^a x) > M\})$. Let $s = C_\mu(\Phi^{a,A}) + \sum_{\gamma \in F} s^a_\gamma$. By hypothesis we have $s < r$. Let $\epsilon > 0$ be small enough so that $s + |F|\epsilon < r$. Since the number $C_\mu(\Phi^{a,A}) = \sum_{\delta \in Q} \mu(A_\delta)$ is independent of $\boldsymbol{a}$, if we can show for each $\gamma \in F$ that the set

$$\{\boldsymbol{b} \in A(\Gamma, X, \mu) \,:\, \mu(\{x \,:\, d_{\Phi^{b,A}}(x, \gamma^b x) > M\}) < s^a_\gamma + \epsilon\} \tag{6.1}$$

contains an open neighborhood of $\boldsymbol{a}$, then the intersection of these sets as γ ranges over F will by Lemma 6.2 be a subset of $A_{F,r}$ containing an open neighborhood of $\boldsymbol{a}$ and we will be done.

Fix then $\gamma \in F$, let $Q^* = Q \cup \{e\}$ and let Σ be the collection

$$\Sigma = \{((\delta_{M-1}, \ldots, \delta_0), (\epsilon_{M-1}, \ldots, \epsilon_0)) \,:\, \delta_j \in Q^* \text{ and } \epsilon_j \in \{-1, 1\} \text{ for } j = 0, \ldots, M-1\}.$$

For each $\boldsymbol{b} \in A(\Gamma, X, \mu)$ and $\sigma \in \Sigma$, writing σ as

$$\sigma = ((\delta_{M-1}, \ldots, \delta_0), (\epsilon_{M-1}, \ldots, \epsilon_0)) \tag{6.2}$$

(where $\delta_j \in Q^*$ and $\epsilon_j \in \{-1, 1\}$ for $j = 0, \ldots, M-1$), we define

$$\varphi^b_\sigma := (\varphi^{b,A}_{\delta_{M-1}})^{\epsilon_{M-1}} \circ \cdots \circ (\varphi^{b,A}_{\delta_0})^{\epsilon_0}.$$

Let $\Sigma(\gamma)$ denote the set of all $\sigma \in \Sigma$ with the property that $\delta_{M-1}^{\epsilon_{M-1}} \cdots \delta_0^{\epsilon_0} = \gamma$. Observe that for $\sigma \in \Sigma(\gamma)$ and $\boldsymbol{b} \in A(\Gamma, X, \mu)$, if $x \in \mathrm{dom}(\varphi^b_\sigma)$ then $\varphi^b_\sigma(x) = \gamma^b x$ and so $d(x, \gamma^b x) \leqslant M$. It follows that

$$\{x \,:\, d_{\Phi^{b,A}}(x, \gamma^b x) > M\} \subseteq \bigcap_{\sigma \in \Sigma(\gamma)} X \setminus \mathrm{dom}(\varphi^b_\sigma). \tag{6.3}$$

If we assume further that $\boldsymbol{b}$ is (essentially) free then, ignoring a null set, the set containment (6.3) becomes an equality. Indeed, restricting to a co-null set X_0 on which b is free we have, for $x \in X_0$, if $d_{\Phi^{b,A}}(x, \gamma^b x) \leqslant M$ then there exists some $\sigma \in \Sigma$ such that $x \in \mathrm{dom}(\varphi^b_\sigma)$ and $\varphi^b_\sigma(x) = \gamma^b x$. Writing σ as in (6.2), this means that $(\delta_{M-1}^{\epsilon_{M-1}} \cdots \delta_0^{\epsilon_0})^b x = \gamma^b x$. Since $\boldsymbol{b}$ is free on X_0 this implies $\delta_{M-1}^{\epsilon_{M-1}} \cdots \delta_0^{\epsilon_0} = \gamma$ and therefore $\sigma \in \Sigma(\gamma)$.

Now, for each $\sigma \in \Sigma$ and $\boldsymbol{b} \in A(\Gamma, X, \mu)$ we see from the definition of φ^b_σ that the set $\mathrm{dom}(\varphi^b_\sigma)$ is an element of the Boolean algebra $\mathcal{A}^b$ generated by

$$\{\alpha^b A_\delta \,:\, \delta \in Q \text{ and } \alpha \in (Q^* \cup Q^{-1})^M\}$$

where $(Q^* \cup Q^{-1})^M = \{\delta_{M-1} \cdots \delta_1 \delta_0 \,:\, \delta_j \in Q^* \cup Q^{-1} \text{ for } j = 0, \ldots, M-1\}$. The algebra $\mathcal{A}^b$ is finite since Q is finite. The Boolean operations are continuous on MALG_μ, so if $\eta > 0$ is small enough (depending on ϵ, Q, and A) then every $\boldsymbol{b}$ in the open neighborhood U_η of $\boldsymbol{a}$ given by

$$U_\eta = \{\boldsymbol{b} \in A(\Gamma, X, \mu) \,:\, \forall \alpha \in (Q^* \cup Q^{-1})^M \, \forall \delta \in Q \ (\mu(\alpha^b A_\delta \Delta \alpha^a A_\delta) < \eta)\}$$

satisfies

$$\mu\Big(\bigcap_{\sigma \in \Sigma(\gamma)} X \setminus \mathrm{dom}(\varphi^b_\sigma)\Big) < \mu\Big(\bigcap_{\sigma \in \Sigma(\gamma)} X \setminus \mathrm{dom}(\varphi^a_\sigma)\Big) + \epsilon = s^a_\gamma + \epsilon$$

where the equality follows from the paragraph following (6.3) since $\boldsymbol{a}$ is free. By (6.3) we then have for such η and $\boldsymbol{b} \in U_\eta$ that

$$\mu(\{x : d_{\Phi^{b},A}(x, \gamma^{b} x) > M\}) < s_\gamma^a + \epsilon$$

which shows that the open neighborhood U_η of $\boldsymbol{a}$ is contained in the set (6.1). □

Note that if $\boldsymbol{a} \in A(\Gamma, X, \mu)$ and $C_\mu(E_a) < r$, then $E = E_a$ witnesses that $\boldsymbol{a}$ satisfies property (1) of Lemma 6.2 and therefore $\boldsymbol{a} \in A_{F,r}(\Gamma, X, \mu)$ for all finite $F \subseteq \Gamma$. It is immediate that if Γ is generated by a finite set F_0 then $A_{F_0,r}(\Gamma, X, \mu) = \{\boldsymbol{a} \in A(\Gamma, X, \mu) : C(\boldsymbol{a}) < r\}$, so we recover (a slightly stronger formulation of) [Ke10, Theorem 10.13] in the following Corollary.

Corollary 6.5 (Kechris, [Ke10]). *Let* Γ *be an infinite, finitely generated group. Then the cost function* $C : A(\Gamma, X, \mu) \to \mathbb{R}$ *is upper semicontinuous at each* $\boldsymbol{a} \in \mathrm{FR}(\Gamma, X, \mu)$*, i.e.,*

$$\limsup_{\boldsymbol{b} \to \boldsymbol{a}} C(\boldsymbol{b}) \leqslant C(\boldsymbol{a}).$$

For general groups, Theorem 6.4 has several consequences for cost and weak containment. It will be helpful to introduce the following notation and definitions.

Definition 6.6. Let $E_0, E_1, E_2, \ldots$, and E be m.p. countable Borel equivalence relations on (X, μ). The sequence $(E_n)_{n\in\mathbb{N}}$ is called an *exhaustion of* E, denoted $(E_n)_{n\in\mathbb{N}} \nearrow E$, if $E_0 \subseteq E_1 \subseteq \cdots$, and $E = \bigcup_n E_n$. The *pseudocost* of E, denoted $PC_\mu(E)$, is defined by

$$PC_\mu(E) = \inf\{\liminf_n C_\mu(E_n) : (E_n)_{n\in\mathbb{N}} \nearrow E\}.$$

If $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ is a m.p. action of a countable group Γ then define the pseudocost of $\boldsymbol{a}$ by $PC(\boldsymbol{a}) := PC_\mu(E_a)$. Finally, define the pseudocost of Γ by $PC(\Gamma) := \inf\{PC(\boldsymbol{a}) : \boldsymbol{a}$ is a free m.p. action of $\Gamma\}$.

It is shown in Corollary 6.17 below that the infimum in the definition of $PC_\mu(E)$ is always attained. If E is aperiodic then $PC_\mu(E) \geqslant 1$ by [KM05, 20.1 and 21.3]. We have $PC_\mu(E) \leqslant C_\mu(E)$ as witnessed by the constant sequence $(E_n)_{n\in\mathbb{N}}$ given by $E_n = E$ for all n. In many cases we actually have the equality $PC_\mu(E) = C_\mu(E)$ as we now show. Recall that a countable Borel equivalence relation E on a standard Borel space X is called *treeable* if there exists an acyclic Borel graph $\mathcal{T} \subseteq X \times X$ whose connected components are the equivalence classes of E. Such a $\mathcal{T}$ is called a *treeing* of E, and we say that E is *treed* by $\mathcal{T}$ to mean that $\mathcal{T}$ is a treeing of E. A theorem of Gaboriau (Theorem 1 of [Ga00]) states that if μ is an E-invariant measure on X and if $\mathcal{T}$ is a treeing of E then $C_\mu(E) = C_\mu(\mathcal{T}) = \frac{1}{2}\int_X \deg_{\mathcal{T}}(x)\, d\mu$. This will be used implicitly below.

Proposition 6.7. *Let* E *be a m.p. countable Borel equivalence relation on* (X, μ) *and let* $(E_n)_{n\in\mathbb{N}}$ *be an exhaustion of* E.

1. *Suppose that* $C_\mu(E) < \infty$*. Then* $C_\mu(E) \leqslant \liminf_n C_\mu(E_n)$.
2. *Suppose that* E *is treeable. Then* $C_\mu(E) \leqslant \liminf_n C_\mu(E_n)$.
3. *(Gaboriau [Ga00]) Suppose that* $\lim_n C_\mu(E_n) = 1$*. Then* $C_\mu(E) = 1$.

In terms of pseudocost vs. cost this implies

Corollary 6.8. *Let* E *be a m.p. countable Borel equivalence relation on* (X, μ).

1. *If* $C_\mu(E) < \infty$ *then* $PC_\mu(E) = C_\mu(E)$.
2. *If* E *is treeable then* $PC_\mu(E) = C_\mu(E)$.

3. $PC_\mu(E) = 1$ *if and only if* $C_\mu(E) = 1$.

Proof of Proposition 6.7. (1): Let $r = \liminf_n C_\mu(E_n)$ and fix $\epsilon > 0$. We may assume that $r < \infty$. Let $\Phi = \{\varphi_i\}_{i=0}^\infty$ be an L-graphing of E with $C_\mu(\Phi) = \sum_{i \geqslant 0} \mu(\mathrm{dom}(\varphi_i)) < \infty$. Let N be so large that $\sum_{i>N} \mu(\mathrm{dom}(\varphi_i)) < \epsilon$. If $M_0 \in \mathbb{N}$ is large enough then for any $n > M_0$ we have $\sum_{i \leqslant N} \mu(\{x \in \mathrm{dom}(\varphi_i) : (x, \varphi_i(x)) \notin E_n\}) < \epsilon$. Since $r = \liminf_n C_\mu(E_n)$ we can find some $n > M_0$ with $C_\mu(E_n) < r + \epsilon$. Let Ψ be an L-graphing of E_n with $C_\mu(\Psi) < r + \epsilon$. Then

$$\Psi \sqcup \{\varphi_i\}_{i>N} \sqcup \{\varphi_i \restriction \{x \in \mathrm{dom}(\varphi_i) : (x, \varphi_i(x)) \notin E_n\}\}_{i \leqslant N}$$

is an L-graphing of E with cost strictly less than $r + 3\epsilon$.

(2): Let $\mathcal{T}$ be a treeing of E and let $\mathcal{T}_n = \mathcal{T} \cap E_n$. Then $\mathcal{T}_n \subseteq \mathcal{T}_{n+1}$ and $\mathcal{T} = \bigcup_n \mathcal{T}_n$ so $\lim_n C_\mu(\mathcal{T}_n) = C_\mu(\mathcal{T})$. Let R_n be the equivalence relation generated by $\mathcal{T}_n$. Then $R_n \subseteq E_n$ and $R_n \cap \mathcal{T} = \mathcal{T}_n$. We need the following lemma which is due to Clinton Conley.

Lemma 6.9 (C. Conley). *Let F be a countable Borel equivalence relation treed by $\mathcal{T}_F$ and let $R \subseteq F$ be a subequivalence relation treed by $\mathcal{T}_R \subseteq \mathcal{T}_F$ (so that $\mathcal{T}_R = R \cap \mathcal{T}_F$). Then any equivalence relation R' with $R \subseteq R' \subseteq F$ has a treeing $\mathcal{T}_{R'}$ with $\mathcal{T}_R \subseteq \mathcal{T}_{R'}$.*

Proof. Proposition 3.3.(iii) of [JKL02] shows how to obtain a treeing $\mathcal{T}_{R'}$ of R' from the given treeing $\mathcal{T}_F$ of F. It is clear from their construction that if an edge of $\mathcal{T}_F$ connects two R'-equivalent points, then that edge remains in $\mathcal{T}_{R'}$. Hence, every edge in $\mathcal{T}_R$ remains in $\mathcal{T}_{R'}$. □[Lemma 6.9]

Apply Lemma 6.9 to $F = E$, $R = R_n$, and $R' = E_n$, along with $\mathcal{T}_F = \mathcal{T}$ and $\mathcal{T}_R = \mathcal{T}_n$, to obtain a treeing $\mathcal{T}'_n$ of E_n with $\mathcal{T}_n \subseteq \mathcal{T}'_n$. Then $\liminf_n C_\mu(E_n) = \liminf_n C_\mu(\mathcal{T}'_n) \geqslant \liminf_n C_\mu(\mathcal{T}_n) = C_\mu(\mathcal{T})$.

(3): Since the E_n are increasing and $\lim_n C_\mu(E_n) = 1$ we have $|[x]_{E_n}| \to \infty$ almost surely (see [KM05, 22.1]), and so E is aperiodic. It follows that $PC_\mu(E) = 1$, so by Corollary 6.17 there is an exhaustion $(E'_n)_{n \in \mathbb{N}}$ of E with $C_\mu(E'_n) \to 1$ such that E'_n is aperiodic for all n. It follows from [KM05, Proposition 23.5] that $C_\mu(E) = 1$. □

Remark 6.10. One may also deduce (2) of Proposition 6.7 by using the equality $C_\mu(E) - 1 = \beta_1(E) - \beta_0(E)$ for treeable E [Ga02, Corollary 3.23] along with [Ga02, Corollary 5.13].

Corollary 6.11. *If E is a m.p. treeable equivalence relation on (X, μ) of infinite cost then any increasing sequence $E_0 \subseteq E_1 \subseteq \cdots$, with $E = \bigcup_n E_n$ satisfies $C_\mu(E_n) \to \infty$.*

Proof. Immediate from (2) of Proposition 6.7. □

Remark 6.12. Corollary 6.11 may be seen as a generalization of a theorem of Takahasi.

Corollary 6.13 (Takahasi [Ta50]). *Suppose $H_0 \subseteq H_1 \subseteq \cdots$ is an ascending chain of subgroups of a free group F, and assume that the H_n have rank uniformly bounded by some natural number $r < \infty$. Then all H_n coincide for n sufficiently large.*

Proof. Suppose that infinitely many H_n are distinct. Then $H = \bigcup_n H_n$ has infinite rank, so Corollary 6.11 implies that for any free m.p. action $H \curvearrowright^a (X, \mu)$ we have $C_\mu(E_{a \restriction H_n}) \to \infty$, contradicting that $\sup_n C_\mu(E_{a \restriction H_n}) \leqslant \sup_n \mathrm{rank}(H_n) \leqslant r$. □

We will use another characterization of pseudocost in order to show that it respects weak containment. In what follows, a sequence $(Q_n)_{n \in \mathbb{N}}$ of subsets of a countable group Γ is called an *exhaustion of* Γ if $Q_0 \subseteq Q_1 \subseteq \cdots$ and $\bigcup_n Q_n = \Gamma$. A sequence $(Q_n)_{n \in \mathbb{N}}$ is called a *finite exhaustion of* Γ if $(Q_n)_{n \in \mathbb{N}}$ is an exhaustion of Γ and Q_n is finite for all $n \in \mathbb{N}$.

Lemma 6.14. *Let E be a m.p. countable Borel equivalence relation on (X, μ) and let $r \in \mathbb{R} \cup \{\infty\}$. Then the following are equivalent:*

1. *There exists an exhaustion $(E_n)_{n \in \mathbb{N}}$ of E with $\limsup_n C_\mu(E_n) \leqslant r$.*
2. *For any countable group Γ and any m.p. action $\boldsymbol{b} = \Gamma \curvearrowright^b (X, \mu)$ with $E = E_b$, and any sequence $(F_n)_{n \in \mathbb{N}}$ of finite subsets of Γ, there exists a finite exhaustion $(Q_n)_{n \in \mathbb{N}}$ of Γ along with an exhaustion $(E_n)_{n \in \mathbb{N}}$ of E such that $F_n \subseteq Q_n$ and $E_{b \restriction \langle Q_n \rangle} \subseteq E_n \subseteq E_{b \restriction \langle Q_{n+1} \rangle}$ for all $n \in \mathbb{N}$, and $\limsup_n C_\mu(E_n) \leqslant r$.*
3. *For any countable group Γ, any m.p. action $\Gamma \curvearrowright^b (X, \mu)$ with $E = E_b$, and any sequence $(F_n)_{n \in \mathbb{N}}$ of finite subsets of Γ, there exists an exhaustion $(E_n)_{n \in \mathbb{N}}$ of E satisfying $E_{b \restriction \langle F_n \rangle} \subseteq E_n$ for all n and $\limsup_n C_\mu(E_n) \leqslant r$.*
4. *For any countable group Γ and any m.p. action $\boldsymbol{b} = \Gamma \curvearrowright^b (X, \mu)$ with $E = E_b$, we have $\boldsymbol{b} \in A_{F, r+\epsilon}$ for all finite $F \subseteq \Gamma$ and all $\epsilon > 0$.*
5. *There exists a countable group Γ and a m.p. action $\boldsymbol{b} = \Gamma \curvearrowright^b (X, \mu)$ with $E = E_b$ such that $\boldsymbol{b} \in A_{F, r+\epsilon}$ for all finite $F \subseteq \Gamma$ and all $\epsilon > 0$.*
6. *There exists a countable group Γ and a m.p. action $\boldsymbol{b} = \Gamma \curvearrowright^b (X, \mu)$ with $E = E_b$, along with an exhaustion $(Q_n)_{n \in \mathbb{N}}$ of Γ and a (not necessarily increasing) sequence $(E_n)_{n \in \mathbb{N}}$ of subequivalence relations of E such that $E_{b \restriction \langle Q_n \rangle} \subseteq E_n$ and $\limsup_n C_\mu(E_n) \leqslant r$.*

Remark 6.15. It is clear that each of the conditions (1), (2), (3), and (6) of Lemma 6.14 are equivalent to their counterparts in which "$\limsup$" is replaced with "$\liminf$" or with "$\lim$."

Proof of 6.14. (1)⇒(4): Assume that $(E_n)_{n \in \mathbb{N}}$ is a sequence as in (1). Let Γ and $\boldsymbol{b} = \Gamma \curvearrowright^b (X, \mu)$ with $E = E_b$ be given. Fix a finite $F \subseteq \Gamma$ and $\epsilon > 0$. Let $n \in \mathbb{N}$ be large enough so that $C_\mu(E_n) < r + \epsilon/2$ and $\sum_{\gamma \in F} \mu(\{x : \gamma^b x \notin [x]_{E_n}\}) < \epsilon/2$. Let $\Phi = \{\gamma^b \restriction \{x : \gamma^b x \notin [x]_{E_n}\}\}_{\gamma \in F}$. Then $R := E_n \vee E_\Phi$ is a subequivalence relation of E containing $E_{b \restriction \langle F \rangle}$ with $C_\mu(R) \leqslant C_\mu(E_n) + C_\mu(\Phi) < r + \epsilon/2 + \epsilon/2 = r + \epsilon$. Then R witnesses that $\boldsymbol{b} \in A_{F, r+\epsilon}(\Gamma, X, \mu)$. This shows that (4) holds.

(4)⇒(2): Assume (4) holds. Let Γ and $\boldsymbol{b} = \Gamma \curvearrowright^b (X, \mu)$ with $E = E_b$ be given along with a sequence $(F_n)_{n \in \mathbb{N}}$ of finite subsets of Γ. We may assume without loss of generality that $(F_n)_{n \in \mathbb{N}}$ is a finite exhaustion of Γ. Fix some sequence of real numbers $\epsilon_n > 0$ with $\epsilon_n \to 0$. We proceed by induction to construct sequences $(Q_n)_{n \in \mathbb{N}}$ and $(E_n)_{n \in \mathbb{N}}$ as in (2). Define $Q_0 = F_0$. Suppose for induction that we have constructed finite subsets $Q_0 \subseteq Q_1 \subseteq \cdots Q_k$ of Γ and equivalence relations $E_0, \dots, E_{k-1}$ with $F_i \subseteq Q_i$ for all $i \leqslant k$ and $E_{b \restriction \langle Q_i \rangle} \subseteq E_i \subseteq E_{b \restriction \langle Q_{i+1} \rangle}$ for all $i < k$. By (4) we have $\boldsymbol{b} \in A_{Q_k \cup F_{k+1}, r+\epsilon_k}$, so by Lemma 6.2 there exists a finite $Q_{k+1} \subseteq \Gamma$ containing $Q_k \cup F_{k+1}$ and a subequivalence relation E_k of E_b with $E_{b \restriction \langle Q_k \rangle} \subseteq E_k \subseteq E_{b \restriction \langle Q_{k+1} \rangle}$ and $C_\mu(E_k) < r + \epsilon_k$. Then Q_{k+1} and E_k extend the induction to the next stage. We obtain from this inductive procedure sequences (Q_n) and (E_n) which satisfy (2) by construction.

(2)⇒(3) is clear. (3)⇒(6) holds since there always exists some countable group Γ and some m.p. action $\boldsymbol{b} = \Gamma \curvearrowright^b (X, \mu)$ with $E = E_b$ (see [FM77]). (6)⇒(5) is routine. Finally, the proof of (4) ⇒ (2) shows that (5) ⇒ (1). □

Remark 6.16. If the the equivalence relation E in Lemma 6.14 is aperiodic then condition (1) implies the stronger statement (1*) in which the equivalence relations E_n are additionally required to be aperiodic. Indeed, assume that E is aperiodic and that (1) holds. Then (3) holds as well. By [Ke10, 3.5] there is an aperiodic $T \in [E]$. Take any countable subgroup $\Gamma \leqslant [E]$ that generates E and with $T \in \Gamma$. Then Γ naturally acts on (X, μ) as a subgroup of $[E]$. Take some finite exhaustion $\{F_n\}_{n \in \mathbb{N}}$ of Γ with $T \in F_0$. Now apply (3) of Lemma 6.14 to this sequence $\{F_n\}_{n \in \mathbb{N}}$ to obtain the desired aperiodic sequence satisfying (1*).

Similarly, if E is aperiodic then (3), and (6) of Lemma 6.14 are each equivalent to their counterparts (3*), and (6*), in which the equivalence relations E_n are each required to be aperiodic.

Corollary 6.17. *Let E be a m.p. countable Borel equivalence relation on (X, μ). There exists an exhaustion $(E_n)_{n\in\mathbb{N}} \nearrow E$ with $\lim_n C_\mu(E_n) = PC_\mu(E)$. In other words, the infimum in the definition of pseudocost is always attained. In addition, if E is aperiodic then such an exhaustion $(E_n)_{n\in\mathbb{N}}$ exists with E_n aperiodic for all n.*

Proof. Let $s = PC_\mu(E)$. By definition of $PC_\mu(E)$, for any $\delta > 0$ there exists a sequence $(E_n^\delta)_{n\in\mathbb{N}} \nearrow E$ with $\limsup_n C_\mu(E_n^\delta) < s + \delta/2$. By [FM77] there is a countable group Γ and some action $\boldsymbol{b} = \Gamma \curvearrowright^b (X, \mu)$ of Γ such that $E = E_b$. Now, E satisfies (1) of Lemma 6.14 with respect to the parameter $r = s + \delta/2$, so by (1)⇒(4) of Lemma 6.14 we have $\boldsymbol{b} \in A_{F,s+\delta/2+\epsilon}$ for all finite $F \subseteq \Gamma$ and $\epsilon > 0$. Taking $\epsilon = \delta/2$ shows that $\boldsymbol{b} \in A_{F,r+\delta}$ for all finite $F \subseteq \Gamma$. Since $\delta > 0$ was arbitrary this shows that $\boldsymbol{b}$ satisfies (5) of Lemma 6.14 with respect to the parameter s, so by (5)⇒(1) Lemma 6.14 there exists a sequence $(E_n)_{n\in\mathbb{N}} \nearrow E$ with $\limsup_n C_\mu(E_n) \leqslant s$. Since $s = PC_\mu(E) \leqslant \liminf_n C_\mu(E_n)$ this shows that in fact $\lim_n C_\mu(E_n) = PC_\mu(E)$. By remark 6.16 if E is aperiodic then we can choose such a sequence $(E_n)_{n\in\mathbb{N}}$ with E_n aperiodic for all n. ☐

Corollary 6.18. *Let E be an aperiodic m.p. countable Borel equivalence relation on (X, μ). Assume that E is ergodic. Then for any exhaustion $(R_n)_{n\in\mathbb{N}}$ of E satisfying $C_\mu(R_n) < \infty$ for all $n \in \mathbb{N}$, there exists an exhaustion $(E_n)_{n\in\mathbb{N}}$ of E with $R_n \subseteq E_n$ for all $n \in \mathbb{N}$ and $\lim_n C_\mu(E_n) = PC_\mu(E)$.*

Proof. Let $(R_n)_{n\in\mathbb{N}}$ be an exhaustion of E with $C_\mu(R_n) < \infty$ for all n. Since E is ergodic we many apply [KM05, Lemma 27.7] to obtain, for each $n \in \mathbb{N}$, a finitely generated group Γ_n and a m.p. action $\boldsymbol{b}_n = \Gamma_n \curvearrowright^{b_n} (X, \mu)$ with $R_n = R_{b_n}$. There is a unique action $\boldsymbol{b} = \Gamma \curvearrowright^a (X, \mu)$ of the free product $\Gamma = *_{n\in\mathbb{N}} \Gamma_n$ satisfying $\boldsymbol{b} \upharpoonright \Gamma_n = \boldsymbol{b}_n$ for all $n \in \mathbb{N}$. For each $n \in \mathbb{N}$ let F_n be a finite generating set for Γ_n. By Corollary 6.17 there exists an exhaustion $(E_n')_{n\in\mathbb{N}}$ of E with $\lim_n C_\mu(E_n') = r$ where $r = PC_\mu(E)$. This shows that E satisfies (1) of Lemma 6.14, so, by applying (3) of Lemma 6.14 to the action $\boldsymbol{b}$ and the sequence $(F_n)_{n\in\mathbb{N}}$, we obtain an exhaustion $(E_n)_{n\in\mathbb{N}}$ of E with $R_n = E_{b\upharpoonright\Gamma_n} \subseteq E_n$ and $\limsup_n C_\mu(E_n) \leqslant r$. Since $r = PC_\mu(E)$ it follows that $\lim_n C_\mu(E_n) = PC_\mu(E)$. ☐

Corollary 6.19. *Let $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ be a m.p. action of Γ. Then $PC(\boldsymbol{a}) \leqslant r$ if and only if $\boldsymbol{a} \in A_{F,r+\epsilon}$ for every finite $F \subseteq \Gamma$ and $\epsilon > 0$.*

Proof. This follows from the equivalence (1)⇔(4) from Lemma 6.14. ☐

Corollary 6.20. *Let $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ and $\boldsymbol{b} = \Gamma \curvearrowright^b (Y, \nu)$ be measure preserving actions of a countable group Γ. Assume that $\boldsymbol{a}$ is free. If $\boldsymbol{a} \prec \boldsymbol{b}$ then $PC(\boldsymbol{b}) \leqslant PC(\boldsymbol{a})$.*

Proof. Let $r = PC(\boldsymbol{a})$. Fix $F \subseteq \Gamma$ finite and $\epsilon > 0$. Since $PC(\boldsymbol{a}) = r$ we have $\boldsymbol{a} \in A_{F,r+\epsilon}(\Gamma, X, \mu)$ by Corollary 6.19. Since $\boldsymbol{a}$ is free, Theorem 6.4 implies that $\boldsymbol{a}$ is contained in the interior of $A_{F,r+\epsilon}(\Gamma, X, \mu)$, so by [Ke10, Proposition 10.1] there exists some $\boldsymbol{c} \in A_{F,r+\epsilon}(\Gamma, X, \mu)$ which is isomorphic to $\boldsymbol{b}$. Hence $\boldsymbol{b} \in A_{F,r+\epsilon}(\Gamma, Y, \nu)$ and therefore $PC(\boldsymbol{b}) \leqslant r$ by Corollary 6.19. ☐

Corollary 6.21. *Let $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ and $\boldsymbol{b} = \Gamma \curvearrowright^b (Y, \nu)$ be measure preserving actions of a countably infinite group Γ. Assume that $\boldsymbol{a}$ is free and is weakly contained in $\boldsymbol{b}$. Then there exists an exhaustion $(E_n)_{n\in\mathbb{N}}$ of E with $\lim_n C_\mu(E_n) \leqslant C(\boldsymbol{a})$ and E_n aperiodic for all $n \in \mathbb{N}$.*

Proof. Corollary 6.20 tells us that $PC(\boldsymbol{b}) \leqslant PC(\boldsymbol{a})$, so by 6.17 we can find an exhaustion $(E_n)_{n\in\mathbb{N}}$ of E, with $\lim_n C_\mu(E_n) \leqslant PC(\boldsymbol{a})$ and E_n aperiodic for all $n \in \mathbb{N}$. Since $PC(\boldsymbol{a}) \leqslant C(\boldsymbol{a})$ we are done. ☐

Corollary 6.22. *Let $\boldsymbol{a}$ and $\boldsymbol{b}$ be m.p. actions of a countably infinite group Γ. Assume that $\boldsymbol{a}$ is free and $\boldsymbol{a} \prec \boldsymbol{b}$.*

1. *If* $\mathrm{C}(\boldsymbol{b}) < \infty$ *then* $\mathrm{C}(\boldsymbol{b}) \leqslant \mathrm{C}(\boldsymbol{a})$.

2. *If* E_b *is treeable then* $\mathrm{C}(\boldsymbol{b}) \leqslant \mathrm{C}(\boldsymbol{a})$.

3. *If* $\mathrm{C}(\boldsymbol{a}) = 1$ *then* $\mathrm{C}(\boldsymbol{b}) = 1$.

Proof. (1) and (2): Suppose $\mathrm{C}(\boldsymbol{b}) < \infty$ or E_b is treeable. Then by Corollary 6.8 and Corollary 6.20 we have $\mathrm{C}(\boldsymbol{b}) = \mathrm{PC}(\boldsymbol{b}) \leqslant \mathrm{PC}(\boldsymbol{a}) \leqslant \mathrm{C}(\boldsymbol{a})$.

Similarly, if $\mathrm{C}(\boldsymbol{a}) = 1$ then by Corollary 6.20 we have $\mathrm{PC}(\boldsymbol{b}) \leqslant \mathrm{PC}(\boldsymbol{a}) \leqslant \mathrm{C}(\boldsymbol{a}) = 1$, so $\mathrm{PC}(\boldsymbol{b}) = 1$ and thus $\mathrm{C}(\boldsymbol{b}) = 1$ by Corollary 6.8. □

Definition 6.23. A group Γ is said to have *fixed price* 1 if $\mathrm{C}(\boldsymbol{a}) = 1$ for every free measure preserving action $\boldsymbol{a}$ of Γ.

In [AW11], Abért and Weiss combine their theorem on free actions (stated above in Theorem 3.1) with [Ke10, Theorem 10.13] to characterize finitely generated groups Γ with fixed price 1 in terms of the Bernoulli shift $\boldsymbol{s}_\Gamma$. We can now remove the hypothesis that Γ is finitely generated.

Corollary 6.24. *Let* Γ *be a countable group. Then the following are equivalent:*

(1) Γ *has fixed price* 1

(2) $\mathrm{C}(\boldsymbol{s}_\Gamma) = 1$

(3) $\mathrm{C}(\boldsymbol{a}) = 1$ *for some m.p. action* $\boldsymbol{a}$ *weakly equivalent to* $\boldsymbol{s}_\Gamma$.

(4) $\mathrm{PC}(\boldsymbol{a}) = 1$ *for some m.p. action* $\boldsymbol{a}$ *weakly equivalent to* $\boldsymbol{s}_\Gamma$.

(5) Γ *is infinite and* $\mathrm{C}(\boldsymbol{a}) \leqslant 1$ *for some non-trivial m.p. action* $\boldsymbol{a}$ *weakly contained in* $\boldsymbol{s}_\Gamma$.

Proof. (1)⇒(2) holds since $\boldsymbol{s}_\Gamma$ is free. (2)⇒(3) is clear. (3) ⇔ (4) follows from Corollary 6.8. Suppose that (3) holds and we will prove (1). Let $\boldsymbol{a}$ be weakly equivalent to $\boldsymbol{s}_\Gamma$ with $\mathrm{C}(\boldsymbol{a}) = 1$. This implies $\boldsymbol{a}$ is free. If $\boldsymbol{b}$ is another free measure preserving action of Γ then $\boldsymbol{a} \prec \boldsymbol{b}$ by Theorem 3.1, so Corollary 6.22 shows that $\mathrm{C}(\boldsymbol{b}) = 1$. Thus Γ has fixed price 1. This shows that properties (1), (2), and (3) are equivalent. The implication (3)⇒(5) is clear.

The proof of the remaining implication (5)⇒(3) uses Lemma 6.34, proved in §6.5 below. Assume that (5) holds. Let $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ be a non-trivial action weakly contained in $\boldsymbol{s}_\Gamma$ with $\mathrm{C}(\boldsymbol{a}) \leqslant 1$. Let $\theta = \theta_{\boldsymbol{a}}$. If Γ is amenable then (1) holds, so we may assume that Γ is non-amenable. Then $\boldsymbol{s}_\Gamma$ is strongly ergodic, hence both $\boldsymbol{a}$ and $\boldsymbol{\theta}$ are weakly mixing. It follows that θ is either a point mass at some finite normal subgroup N of Γ, or θ concentrates on the infinite subgroups of Γ.

Case 1: θ is a point mass at some finite normal subgroup $N \leqslant \Gamma$. Then $\mathrm{C}(\boldsymbol{a}) = 1$ since E_a is aperiodic. By [CKT-D12, Proposition 4.7] there is some $\boldsymbol{b} = \Gamma \curvearrowright^b (Y, \nu)$ weakly equivalent to $\boldsymbol{s}_\Gamma$ such that $\boldsymbol{a}$ is a factor of $\boldsymbol{b}$, say via the factor map $\pi : Y \to X$. Let Y_0 be a Borel transversal for the orbits of $N \curvearrowright^b (Y, \nu)$ and let $\sigma : Y \to Y_0$ be the corresponding selector. Let ν_0 denote the normalized restriction of ν to Y_0 and let $\boldsymbol{b}_0$ be the action of Γ on (Y_0, ν_0) given by $\gamma^{b_0} y = \sigma(\gamma^b y)$. Then π factors $\boldsymbol{b}_0$ onto $\boldsymbol{a}$. Since $\theta_{\boldsymbol{a}} = \theta_{\boldsymbol{b}_0} = \delta_N$, the actions $\boldsymbol{a}$ and $\boldsymbol{b}_0$ descend to free actions $\tilde{\boldsymbol{a}}$ and $\tilde{\boldsymbol{b}}_0$ respectively of Γ/N, and π factors $\tilde{\boldsymbol{b}}_0$ onto $\tilde{\boldsymbol{a}}$. Then $\mathrm{C}(\tilde{\boldsymbol{a}}) = \mathrm{C}(\boldsymbol{a}) = 1$, so $\mathrm{C}(\tilde{\boldsymbol{b}}_0) = 1$ by Corollary 6.22. Since $\mathrm{E}_{b_0} = \mathrm{E}_b \upharpoonright Y_0$ we have $\mathrm{C}_{\nu_0}(\mathrm{E}_b \upharpoonright Y_0) = 1$, so $\mathrm{C}(\boldsymbol{b}) = \mathrm{C}_\nu(\mathrm{E}_b) = 1$ by [KM05, Theorem 25.1] ([KM05, Theorem 21.1] also works). This shows that (3) holds.

Case 2: θ is infinite. We have $\boldsymbol{a} \prec \boldsymbol{s}_\Gamma$, so $\boldsymbol{a}$ is NA-ergodic and therefore θ is amenable by Theorem 3.13. Then $\mathrm{C}(\boldsymbol{\theta}_{\boldsymbol{a}} \times \boldsymbol{s}_\Gamma) = 1$ by Lemma 6.34, and $\boldsymbol{\theta}_{\boldsymbol{a}} \times \boldsymbol{s}_\Gamma$ is weakly equivalent to $\boldsymbol{s}_\Gamma$, so (3) holds. □

Note 6.25. Similar to [Ke10, Corollary 10.14], one may strengthen Corollaries 6.20, 6.21, and 6.22 by replacing the hypothesis $\boldsymbol{a} \prec \boldsymbol{b}$ their statements with the weaker hypothesis that

$$\boldsymbol{a} \in \overline{\{\boldsymbol{c} \in \mathrm{A}(\Gamma, X, \mu) : E_c \text{ is orbit equivalent to } E_b\}} \tag{6.4}$$

where (X, μ) is the underlying space of $\boldsymbol{a}$. The proofs remain the same. Note that (6.4) is actually slightly weaker than the hypothesis $\boldsymbol{a} \preceq \boldsymbol{b}$ from [Ke10, Corollary 10.14], since the action $\boldsymbol{c}$ from (6.4) ranges over all of $\mathrm{A}(\Gamma, X, \mu)$ and not just $\mathrm{FR}(\Gamma, X, \mu)$. Specializing to the case where Γ is finitely generated, we recover a somewhat strengthened version of the first statement of [Ke10, Corollary 10.14].

6.3 The cost of a generic action

The results of the previous section have consequences for generic properties (with respect to the weak topology) in $\mathrm{FR}(\Gamma, X, \mu)$ related to cost. We begin by proving analogues of Corollaries 6.17 and 6.8 for groups. Recall that a countable group Γ is called *treeable* if it admits a free measure preserving action $\boldsymbol{a}$ such that E_a is treeable.

Proposition 6.26. *Let Γ be a countably infinite group.*

(1) Suppose that $C(\Gamma) < \infty$. Then for any free m.p. action $\boldsymbol{b} = \Gamma \curvearrowright^b (X, \mu)$ of Γ, and any exhaustion $(E_n)_{n \in \mathbb{N}}$ of E_b, we have $\liminf_{n\to\infty} C_\mu(E_n) \geqslant C(\Gamma)$. Hence $\mathrm{PC}(\Gamma) = C(\Gamma)$.

(2) Suppose that Γ is treable. Then $\mathrm{PC}(\Gamma) = C(\Gamma)$.

(3) $\mathrm{PC}(\Gamma) = 1$ if and only if $C(\Gamma) = 1$.

(4) $\mathrm{PC}(\Gamma)$ is attained by some free m.p. action of Γ. In fact, if $\boldsymbol{a} \in FR(\Gamma, X, \mu)$ has dense conjugacy class in $(FR(\Gamma, X, \mu), w)$ then $\mathrm{PC}(\boldsymbol{a}) = \mathrm{PC}(\Gamma)$.

Proof. (1): Let $\boldsymbol{b}$ be a free m.p. action of Γ. It suffices to show that $\mathrm{PC}(\boldsymbol{b}) \geqslant C(\Gamma)$. Let $\boldsymbol{a}$ be a free m.p. action of Γ with $C(\boldsymbol{a}) = C(\Gamma) < \infty$ and let $\boldsymbol{c} = \boldsymbol{a} \times \boldsymbol{b}$. Then by the remark at the bottom of p. 78 in [Ke10] we have $C(\boldsymbol{c}) \leqslant C(\boldsymbol{a}) = C(\Gamma)$, hence $C(\boldsymbol{c}) = C(\Gamma) < \infty$. Since $C(\boldsymbol{c}) < \infty$ we have $\mathrm{PC}(\boldsymbol{c}) = C(\boldsymbol{c})$ by (1) of Corollary 6.8. In addition, $\boldsymbol{b} \prec \boldsymbol{c}$ and $\boldsymbol{b}$ is free, so Corollary 6.20 implies $\mathrm{PC}(\boldsymbol{b}) \geqslant \mathrm{PC}(\boldsymbol{c}) = C(\boldsymbol{c}) = C(\Gamma)$.

(2): Let $\boldsymbol{b}$ be a free m.p. action of Γ. Once again it suffices to show $\mathrm{PC}(\boldsymbol{b}) \geqslant C(\Gamma)$. Let $\boldsymbol{a}$ be a free m.p. action of Γ with E_a treeable and let $\boldsymbol{c} = \boldsymbol{a} \times \boldsymbol{b}$. By [KM05, Proposition 30.5] E_c is treeable and $C(\boldsymbol{c}) = C(\boldsymbol{a}) = C(\Gamma)$. Then (2) of Corollary 6.8 implies that $\mathrm{PC}(\boldsymbol{c}) = C(\boldsymbol{c})$, so, as $\boldsymbol{b} \prec \boldsymbol{c}$, Corollary 6.20 implies that $\mathrm{PC}(\boldsymbol{b}) \geqslant \mathrm{PC}(\boldsymbol{c}) = C(\boldsymbol{c}) = C(\Gamma)$.

(3): This is immediate from (3) of Corollary 6.8.

(4): If $\boldsymbol{a} \in \mathrm{FR}(\Gamma, X, \mu)$ has dense conjugacy class this means that $\boldsymbol{b} \prec \boldsymbol{a}$ for every m.p. action $\boldsymbol{b}$ of Γ [Ke10, Proposition 10.1] (also note that such an $\boldsymbol{a}$ exists by [Ke10, Theorem 10.7]). Corollary 6.20 then shows that $\mathrm{PC}(\boldsymbol{a}) \leqslant \inf\{\mathrm{PC}(\boldsymbol{b}) : \boldsymbol{b} \in \mathrm{FR}(\Gamma, X, \mu)\} = \mathrm{PC}(\Gamma)$, hence $\mathrm{PC}(\boldsymbol{a}) = \mathrm{PC}(\Gamma)$. ☐

By [Ke10, Proposition 10.10] the cost function $\boldsymbol{a} \mapsto C(\boldsymbol{a})$ is constant on a dense G_δ subset of $\mathrm{FR}(\Gamma, X, \mu)$. Let $C_{\mathrm{gen}}(\Gamma) \in [0, \infty]$ denote this constant value. Similarly, the pseudocost function $\boldsymbol{a} \mapsto \mathrm{PC}(\boldsymbol{a})$ is constant on a dense G_δ subset of $\mathrm{FR}(\Gamma, X, \mu)$. Denote this constant value by $\mathrm{PC}_{\mathrm{gen}}(\Gamma)$. Problem 10.11 of [Ke10] asks whether $C_{\mathrm{gen}}(\Gamma) = C(\Gamma)$ holds for every countably infinite group Γ, and [Ke10, Corollary 10.14] shows that the equality holds whenever Γ is finitely generated.

Corollary 6.27. *Let Γ be a countably infinite group. Then*

1. *The set $\mathrm{MINPCOST}(\Gamma, X, \mu) = \{\boldsymbol{a} \in \mathrm{FR}(\Gamma, X, \mu) : \mathrm{PC}(\boldsymbol{a}) = \mathrm{PC}(\Gamma)\}$ is dense G_δ in $\mathrm{A}(\Gamma, X, \mu)$. In particular, $\mathrm{PC}_{gen}(\Gamma) = \mathrm{PC}(\Gamma)$.*

2. *Either* $C_{gen}(\Gamma) = C(\Gamma)$ *or* $C_{gen}(\Gamma) = \infty$.

3. *If* $PC(\Gamma) = 1$ *then* $C_{gen}(\Gamma) = C(\Gamma) = 1$.

Proof. (1): Let $r = PC(\Gamma)$. Corollary 6.19 shows that

$$\mathrm{MINPCOST}(\Gamma, X, \mu) = \bigcap \{A_{F,r+1/n}(\Gamma, X, \mu) \cap \mathrm{FR}(\Gamma, X, \mu) \,:\, F \subseteq \Gamma \text{ is finite and } n \in \mathbb{N}\}.$$

To show this set is dense G_δ in $A(\Gamma, X, \mu)$ it therefore suffices to show that $A_{F,r+\epsilon}(\Gamma, X, \mu) \cap \mathrm{FR}(\Gamma, X, \mu)$ is dense G_δ for each $F \subseteq \Gamma$ finite and $\epsilon > 0$. By [Ke10, Theorem 10.8], the set $\mathrm{FR}(\Gamma, X, \mu)$ is dense G_δ in $A(\Gamma, X, \mu)$. Theorem 6.4 shows that $A_{F,r+\epsilon}$ is relatively open in $\mathrm{FR}(\Gamma, X, \mu)$, so it only remains to show that it is dense. By Proposition 6.26 we have $PC(\boldsymbol{a}) = PC(\Gamma)$ whenever $\boldsymbol{a} \in \mathrm{FR}(\Gamma, X, \mu)$ has a dense conjugacy class. Since the set of actions with dense conjugacy class is dense G_δ in $\mathrm{FR}(\Gamma, X, \mu)$ the result follows.

(2): Suppose that $C_{\mathrm{gen}}(\Gamma) = r < \infty$. This means the generic $\boldsymbol{a} \in \mathrm{FR}(\Gamma, X, \mu)$ has $C(\boldsymbol{a}) = r$. Since $r < \infty$ it follows from Corollary 6.8 that $C(\boldsymbol{a}) = r \Rightarrow C(\boldsymbol{a}) = PC(\boldsymbol{a})$. Thus the generic free action $\boldsymbol{a}$ satisfies $PC(\boldsymbol{a}) = r = C(\boldsymbol{a})$ and by part (1) we therefore have $C(\Gamma) \geqslant PC(\Gamma) = PC_{\mathrm{gen}} = C_{\mathrm{gen}}(\Gamma) \geqslant C(\Gamma)$, which shows that $C_{\mathrm{gen}}(\Gamma) = C(\Gamma)$.

(3) follows from (1) along with Corollary 6.8. ☐

Let $\mathrm{MINCOST}(\Gamma, X, \mu) = \{\boldsymbol{a} \in \mathrm{FR}(\Gamma, X, \mu) \,:\, C(\boldsymbol{a}) = C(\Gamma)\}$.

Corollary 6.28. *Let* Γ *be a countably infinite group. Then the set*

$$D = \Big\{\boldsymbol{b} \in \mathrm{FR}(\Gamma, X, \mu) : \exists \textit{aperiodic subequivalence relations } E_0 \subseteq E_1 \subseteq E_2 \subseteq \cdots \textit{ of } E_{\boldsymbol{b}}, \textit{ with } E_{\boldsymbol{b}} = \bigcup_n E_n \textit{ and } \lim_n C_\mu(E_n) = C(\Gamma)\Big\}$$

is dense G_δ *in* $A(\Gamma, X, \mu)$. *Additionally, if* $C(\Gamma) < \infty$ *then we have the equality of sets*

$$\mathrm{MINCOST}(\Gamma, X, \mu) = D \cap \{\boldsymbol{b} \in \mathrm{FR}(\Gamma, X, \mu) \,:\, C(\boldsymbol{b}) < \infty\}. \tag{6.5}$$

In particular, if all free actions of Γ *have finite cost then* $\mathrm{MINCOST}(\Gamma, X, \mu) = D$ *is dense* G_δ.

Proof. We begin by showing D is dense G_δ. By [Ke10, Theorem 10.8], $\mathrm{FR}(\Gamma, X, \mu)$ is dense G_δ in $A(\Gamma, X, \mu)$. If $C(\Gamma) = \infty$ then $D = \mathrm{FR}(\Gamma, X, \mu)$ and we are done, so we may assume that $C(\Gamma) < \infty$. Then $C(\Gamma) = PC(\Gamma)$ by Proposition 6.26, so it follows from Corollary 6.17 that $D = \{\boldsymbol{a} \in \mathrm{FR}(\Gamma, X, \mu) \,:\, PC(\boldsymbol{a}) = PC(\Gamma)\} = \mathrm{MINPCOST}(\Gamma, X, \mu)$, and therefore D is dense G_δ by Corollary 6.27.

For the second statement of the theorem, suppose that $C(\Gamma) < \infty$. Then $C(\Gamma) = PC(\Gamma)$ by Proposition 6.26. The inclusion from left to right in (6.5) is clear. If $\boldsymbol{b}$ has finite cost and $\boldsymbol{b} \in D$ then, $PC(\boldsymbol{b}) \leqslant C(\Gamma) = PC(\Gamma)$, hence $PC(\boldsymbol{b}) = PC(\Gamma) = C(\Gamma)$, i.e., $\boldsymbol{b} \in \mathrm{MINCOST}(\Gamma, X, \mu)$. ☐

6.4 Cost and invariant random subgroups

Equip each of the spaces Γ^Γ and 2^Γ with the pointwise convergence topology.

Lemma 6.29. *There exists a continuous assignment* $\mathrm{Sub}_\Gamma \to \Gamma^\Gamma$, $H \mapsto \sigma_H$, *with the following properties:*

(i) *For each* $H \in \mathrm{Sub}_\Gamma$, $\sigma_H : \Gamma \to \Gamma$ *is a selector for the right cosets of* H *in* Γ, *i.e.,* $\sigma_H(\delta) \in H\delta$ *for all* $\delta \in \Gamma$, *and* σ_H *is constant on each right coset of* H.

(ii) $\sigma_H(h) = e$ *whenever* $h \in H$.

(iii) The corresponding assignment of transversals $\mathrm{Sub}_\Gamma \to 2^\Gamma$, $H \mapsto T_H := \sigma_H(\Gamma)$, *is continuous.*

Proof. Fix a bijective enumeration $\Gamma = \{\gamma_m\}_{m\in\mathbb{N}}$ of Γ with $\gamma_0 = e$, and define $\sigma_H(\gamma_m) = \gamma_i$ where i is least such that $\gamma_m\gamma_i^{-1} \in H$. This is continuous and (i) and (ii) are clearly satisfied, and (iii) follows from continuity of $H \mapsto \sigma_H$, since the map $\Gamma^\Gamma \to 2^\Gamma$ sending $f : \Gamma \to \Gamma$ to its set of fixed points is continuous. ☐

Define the set

$$A(\mathrm{Sub}_\Gamma, X, \mu) := \{(H, \boldsymbol{a}) : H \in \mathrm{Sub}_\Gamma, \text{ and } \boldsymbol{a} \in A(H, X, \mu)\}.$$

This set has a natural Polish topology in which $(H_n, \boldsymbol{a}_n) \to (H, \boldsymbol{a})$ if and only if $H_n \to H$ and $\boldsymbol{a}_n \to \boldsymbol{a}$ pointwise. We make this precise by taking $*$ to be some point isolated from $\mathrm{Aut}(X, \mu)$ and then defining $\gamma^b = *$ whenever $H \leqslant \Gamma$, $\boldsymbol{b} \in A(H, X, \mu)$, and $\gamma \notin H$. Then $(H_n, \boldsymbol{a}_n) \to (H, \boldsymbol{a})$ means that $\gamma^{a_n} \to \gamma^a$ for every $\gamma \in \Gamma$.

Lemma 6.30. *For any* $r \in \mathbb{R}$ *the sets*

$$S_r = \{H \in \mathrm{Sub}_\Gamma : C(H) < r\}$$
$$A_r = \{(H, \boldsymbol{a}) \in A(\mathrm{Sub}_\Gamma, X, \mu) : \boldsymbol{a} \text{ is free and } C(\boldsymbol{a}) < r\}$$

are analytic. In particular, the map $H \mapsto C(H)$ *is universally measurable.*

Proof. It suffices to show that A_r is analytic since S_r is the image of A_r under projection onto Sub_Γ which is continuous. We may assume that $X = 2^\mathbb{N}$ and that μ is the uniform product measure.

Let $\Gamma \curvearrowright^s X^\Gamma$ denote the left shift action given by $(\gamma^s \cdot f)(\delta) = f(\gamma^{-1}\delta)$ for $f \in X^\Gamma$. Let $H \mapsto \sigma_H$ and $H \mapsto T_H \subseteq \Gamma$ be a continuous assignment of selectors and transversals given by Lemma 6.29. For $(H, \boldsymbol{a}) \in A(\mathrm{Sub}_\Gamma, X, \mu)$ define the map $\Phi_{H,\boldsymbol{a}} : X \to X^\Gamma$ by $\Phi_{H,\boldsymbol{a}}(x)(ht) = (h^{-1})^a x$ for $h \in H$, $t \in T_H$, $x \in X$. Then $\Phi_{H,\boldsymbol{a}}$ is injective and equivariant from $H \curvearrowright^a X$ to the shift action $H \curvearrowright^s X^\Gamma$ and so the measure $\mu_{H,\boldsymbol{a}} := (\Phi_{H,\boldsymbol{a}})_*\mu$ is $H \curvearrowright^s X^\Gamma$ invariant, and the systems $H \curvearrowright^a (X, \mu)$ and $H \curvearrowright^s (X^\Gamma, \mu_{H,\boldsymbol{a}})$ are isomorphic. Let P denote the space of Borel probability measures on X^Γ equipped with the weak*-topology.

Claim 1. *The map* $A(\mathrm{Sub}_\Gamma, X, \mu) \to P$, $(H, \boldsymbol{a}) \mapsto \mu_{H,\boldsymbol{a}}$ *is continuous.*

Proof of claim. Suppose that $(H_n, \boldsymbol{a}_n) \to (H_\infty, \boldsymbol{a}_\infty)$ in $A(\mathrm{Sub}_\Gamma, X, \mu)$. Letting $\mu_n = \mu_{H_n,\boldsymbol{a}_n}$, it suffices to check that $\mu_n(A) \to \mu_\infty(A)$ whenever $A \subseteq X^\Gamma$ is of the form $A = \{f \in X^\Gamma : \forall\gamma \in F\ (f(\gamma) \in A_\gamma)\}$ where $F \subseteq \Gamma$ is finite and $A_\gamma \subseteq X$ is Borel. For $\gamma \in F$ write $\gamma = h_\gamma t_\gamma$ where $t_\gamma \in T_{H_\infty}$ and $h_\gamma \in H_\infty$. By continuity of $H \mapsto \sigma_H$ and $H \mapsto T_H$, for all large enough n, $h_\gamma \in H_n$ and $t_\gamma \in T_{H_n}$ for all $\gamma \in F$. Then $\mu_n(A) = \mu(\bigcap_{\gamma\in F} h_\gamma^{a_n}(A_\gamma)) \to \mu(\bigcap_{\gamma\in F} h_\gamma^{a}(A_\gamma)) = \mu_\infty(A)$ since $\boldsymbol{a}_n \to \boldsymbol{a}$. ☐[Claim]

Now let E_H denote the orbit equivalence relation on X^Γ generated by $H \curvearrowright^s X^\Gamma$. The set

$$B = \{(H, \nu) \in \mathrm{Sub}_\Gamma \times P : \nu \text{ is } E_H\text{-invariant and } H \curvearrowright^s (X^\Gamma, \nu) \text{ is essentially free}\}$$

is Borel so by the proof of [KM05, Proposition 18.1] the set $D = \{(H, \nu) \in B : C_\nu(E_H) < r\}$ is analytic. We have $(H, \boldsymbol{a}) \in A_r$ if and only if $(H, \mu_{H,\boldsymbol{a}}) \in D$, which shows that A_r is analytic. ☐

It follows that for any ergodic invariant random subgroup θ of Γ there is an $r \in \mathbb{R} \cup \{\infty\}$ such that $C(H) = r$ for almost all $H \leqslant \Gamma$. The following is an analogue of [BG05, §5] for cost. I would like to thank Lewis Bowen for a helpful discussion related to this.

Theorem 6.31. *Let* θ *be an invariant random subgroup of* Γ *and suppose that* θ *concentrates on the infinite subgroups of* Γ *which have infinite index in* Γ. *If* $\theta(\{H : C(H) < \infty\}) \neq 0$ *then* $C(\Gamma) = 1$.

Thus, if $C(\Gamma) > 1$ *then for any ergodic non-atomic m.p. action* $\Gamma \curvearrowright^a (X, \mu)$, *either* Γ_x *is finite almost surely, or* $C(\Gamma_x) = \infty$ *almost surely.*

Proof. To see that the second statement follows from the first observe that an ergodic non-atomic m.p. action cannot have stabilizers which are finite index. We now prove the first statement. By decomposing θ into its ergodic components we may assume without loss of generality that θ is ergodic and there is an $r \in \mathbb{R}$ such that $C(H) < r$ almost surely.

By Lemma 6.30 the set $A_r = \{(H, \boldsymbol{a}) \in A(\mathrm{Sub}_\Gamma, X, \mu) : \boldsymbol{a}$ is free and $C(\boldsymbol{a}) < r\}$ is an analytic subset of $A(\mathrm{Sub}_\Gamma, X, \mu)$. Since $C(H) < r$ almost surely, we may measurably select for each $H \in \mathrm{Sub}_\Gamma$ a free action $\boldsymbol{a}_H \in \mathrm{FR}(H, X, \mu) \subseteq A(H, X, \mu)$ of H such that almost surely $C(\boldsymbol{a}_H) < r$ (we are applying [Ke95, 18.1] to the flip of the graph of the projection function $A_r \to \mathrm{Sub}_\Gamma$, $(H, \boldsymbol{a}) \mapsto H$). A co-inducing process can now be used to obtain an action $\boldsymbol{b}$ of Γ from the selection $H \mapsto \boldsymbol{a}_H \in A(H, X, \mu)$ as follows.

Let $H \mapsto \sigma_H$ be as in Lemma 6.29. Let $\mathrm{COS}_\Gamma \subseteq 2^\Gamma$ denote the closed subspace of all right cosets of subgroups of Γ, on which Γ acts continuously by left translation $\gamma^\ell \cdot H\delta = \gamma H\delta$. The function $\rho : \Gamma \times \mathrm{COS}_\Gamma \to \Gamma$ defined by

$$\rho(\gamma, H\delta) = (\sigma_{\gamma H\gamma^{-1}}(\gamma\delta))^{-1}\gamma\sigma_H(\delta)$$

is a continuous cocycle of this action with values in Γ. It is clear that $\rho(\gamma, H\delta) \in \delta^{-1}H\delta$, so the map $(\gamma, H\delta) \mapsto \rho(\gamma, H\delta)^{a_{\delta^{-1}H\delta}}$ is a well-defined measurable cocycle with values in $\mathrm{Aut}(X, \mu)$. We therefore obtain an action b of Γ on the space $W = \{(H, f) : H \leqslant \Gamma$ and $f : H\backslash\Gamma \to X\}$ given by $\gamma^b(H, f) = (\gamma H\gamma^{-1}, \gamma^{b_H} f)$ where $\gamma^{b_H} f : \gamma H\gamma^{-1}\backslash\Gamma \to X$ is given by

$$(\gamma^{b_H} f)(\gamma H\delta) = \rho(\gamma, H\delta)^{a_{\delta^{-1}H\delta}}(f(H\delta)).$$

This action preserves the measure $\kappa = \int_H (\delta_H \times \mu^{H\backslash\Gamma})\, d\theta(H)$ since

$$\begin{aligned}\gamma^b_*\kappa &= \int_H (\delta_{\gamma H\gamma^{-1}} \times \gamma^{b_H}_* \mu^{H\backslash\Gamma})\, d\theta = \int_H \Big(\delta_{\gamma H\gamma^{-1}} \times \prod_{\gamma H\delta \in \gamma H\gamma^{-1}\backslash\Gamma} (\rho(\gamma, H\delta)^{a_{\delta^{-1}H\delta}})_*\mu\Big)\, d\theta \\ &= \int_H \Big(\delta_{\gamma H\gamma^{-1}} \times \prod_{\gamma H\delta \in \gamma H\gamma^{-1}\backslash\Gamma} \mu\Big)\, d\theta = \int_H (\delta_{\gamma H\gamma^{-1}} \times \mu^{\gamma H\gamma^{-1}\backslash\Gamma})\, d\theta = \int_H \delta_H \times \mu^{H\backslash\Gamma} d\theta = \kappa.\end{aligned}$$

Lemma 6.32.

1. *For each* $(H, f) \in W$, *and* $h \in H$ *we have* $(h^{b_H} f)(H) = h^{a_H}(f(H))$ *and thus the map* $X^{H\backslash\Gamma} \to X$, $f \mapsto f(H)$ *factors*

$$\boldsymbol{b}_H = H \curvearrowright^{b_H} (X^{H\backslash\Gamma}, \mu^{H\backslash\Gamma})$$

onto $\boldsymbol{a}_H$.

2. *(Analogue of [Io11, Lemma 2.1]) For almost all* $H \leqslant \Gamma$ *and every* $\gamma \in \Gamma \setminus \{e\}$ *the sets*

$$W^H_\gamma = \{f \in X^{H\backslash\Gamma} : \gamma H\gamma^{-1} = H \textit{ and } (\gamma^{b_H} f)(H) = f(H)\}$$

are $\mu^{H\backslash\Gamma}$*-null. In particular,* $\boldsymbol{b}$ *is essentially free.*

Proof. (1) is clear from the definition of b_H. For (2), If $f \in W^H_\gamma$ then $\rho(\gamma, H)^{a_H}(f(H\gamma^{-1})) = f(H)$ by definition of b_H. So for each H with $\boldsymbol{a}_H$ essentially free, if $\gamma \in H \setminus \{e\}$ then $f \in W^H_\gamma$ if and only if $\gamma^{a_H}(f(H)) = f(H)$, so that W^H_γ is null, while if $\gamma \in \Gamma \setminus H$ then $W^H_\gamma \subseteq \{f \in X^{H\backslash\Gamma} : \rho(\gamma, H)^{a_H}(f(H\gamma^{-1})) = f(H)\}$, which is null since $H\gamma^{-1} \neq H$ and μ is non-atomic. Since almost all $\boldsymbol{a}_H$ are essentially free we are done. ☐[Lemma 6.32]

We now apply a randomized version of an argument due to Gaboriau (see [KM05, Theorem 35.5]). There is another measure preserving action $\boldsymbol{s} = \Gamma \curvearrowright^s (W, \kappa)$ of Γ on (W, κ) given by $\gamma^s(H, f) = (\gamma H\gamma^{-1}, \gamma^{s_H} f)$ where $(\gamma^{s_H} f)(\gamma H\delta) = f(H\delta)$ (this is the random Bernoulli shift determined by θ [T-D12a, §5.3]). The projection map $W \to \mathrm{Sub}_\Gamma$, $(H, f) \mapsto H$ factors both $\boldsymbol{b}$ and $\boldsymbol{s}$ onto $\boldsymbol{\theta}$. We let $\boldsymbol{a}$ denote the corresponding relatively independent joining of $\boldsymbol{b}$ and $\boldsymbol{s}$ over $\boldsymbol{\theta}$, i.e., $\boldsymbol{a}$ is the measure preserving action of Γ on

$$(Z, \eta) = \Big(\big\{(H, f, g) \,:\, f, g \in X^{H\backslash\Gamma}\big\},\ \int_H (\delta_H \times \mu^{\Gamma/H} \times \mu^{\Gamma/H})\, d\theta\Big)$$

given by $\gamma^a(H, f, g) = (\gamma H \gamma^{-1}, \gamma^{b_H} f, \gamma^{s_H} g)$ where $(\gamma^{s_H} g)(\gamma H\delta) = g(H\delta)$. This action is free since it factors onto $\boldsymbol{b}$.

Let $p : Z \to W$ denote the projection map $p((H, f, g)) = (H, g)$. For each $(H, g) \in W$ the set $p^{-1}((H, g))$ is $a \restriction H$-invariant, and we let $E_{(H,g)}$ denote the orbit equivalence relation on $p^{-1}((H, g))$ generated by $\boldsymbol{a} \restriction H$, i.e., $(H, f_1, g)E_{(H,g)}(H, f_2, g)$ if and only if there is some $h \in H$ such that $h^{b_H} f_1 = f_2$. Define the equivalence relation E on Z by $E = \bigsqcup_{(H,g)\in W} E_{(H,g)}$, i.e.,

$$(H_1, f_1, g_1)E(H_2, f_2, g_2) \Leftrightarrow (H_1, g_1) = (H_2, g_2) \text{ and } \exists h \in H_1\ (h^{b_H} f_1 = f_2).$$

Recall that if $F \subseteq R$ are countable Borel equivalence relations on a standard Borel space Y, then F is said to be *normal* in R if there exists some countable group Δ of Borel automorphisms of Y which generates R and satisfies $xFy \Rightarrow \delta(x)F\delta(y)$ for all $\delta \in \Delta$.

Lemma 6.33. *E is a normal subequivalence relation of E_a that is almost everywhere aperiodic and with $C_\eta(E) < r$.*

Proof of Lemma 6.33. It is clear that E is an equivalence relation and that E is contained in E_a. Also, E is almost everywhere aperiodic since θ concentrates on the infinite subgroups of Γ by hypothesis. Let $\gamma \in \Gamma$ and let $(H, f, g), (H, f', g) \in \mathrm{Sub}_\Gamma \times X$ be E-related so that $h^{b_H} f = f'$ for some $h \in H$. To show E is normal in E_a we must show that $\gamma^a(H, f, g)$ and $\gamma^a(H, h^{b_H} f, g)$ are E related as well, i.e., we must find some $k \in \gamma H \gamma^{-1}$ such that $(k\gamma)^{b_H} f_1 = \gamma^{b_H}(h^{b_H} f_1)$. The element $k = \gamma h \gamma^{-1}$ works.

If we disintegrate η via the E-invariant map $p : Z \to W$, then for each $(H, g) \leqslant \Gamma$, the equivalence relation $E_{(H,g)}$ on $(p^{-1}((H, g)), \eta_{(H,g)})$ is isomorphic to the orbit equivalence relation generated by $b_H \restriction H$ on $(X^{H\backslash\Gamma}, \mu^{H\backslash\Gamma})$. By Lemma 6.32.(1), $\boldsymbol{b}_H$ factors onto $\boldsymbol{a}_H$, so for θ-almost every H we have $r \leqslant C_{\eta_{(H,g)}}(E_{(H,g)}) = C(\boldsymbol{b}_H) \leqslant C(\boldsymbol{a}_H) < r$ by [Ke10, bottom of p. 78]. Then by [KM05, Proposition 18.4] we have

$$C_\eta(E) = \int_{H,g} C_{\eta_{(H,g)}}(E_{(H,g)})\, d\theta(H) < r. \qquad \square\text{[Lemma 6.33]}$$

Since H is almost surely infinite index, the equivalence relation E_s on W generated by $\boldsymbol{s}$ is aperiodic. By [Ke10] the full group $[E_s]$ contains an aperiodic transformation $T : W \to W$. Let $B : \Gamma \to \mathrm{MALG}_\kappa$, $\gamma \mapsto B_\gamma$, be a partition of W such that $T \restriction B_\gamma = \gamma^s \restriction B_\gamma$. Then $A : \Gamma \to \mathrm{MALG}_\kappa$ given by $A_\gamma = p^{-1}(B_\gamma)$ is a partition of Z, and determines the L-graphing $\Phi^{a,A} = \{\varphi^{a,A}_\gamma\}_{\gamma\in\Gamma}$ where $\varphi^{a,A}_\gamma \restriction A_\gamma = \gamma^a \restriction A_\gamma$.

Fix $\epsilon > 0$ and find by Lemma 6.33 a graphing $\{\varphi_i\}_{i\in\mathbb{N}}$ of $E \subseteq Z$ of finite cost $\sum_i C_\eta(\varphi_i) < \infty$. Let M be so large that $\sum_{i>M} C_\eta(\varphi_i) < \epsilon/2$. Let $Y_0 \subseteq W$ be a Borel complete section for E_T with $\kappa(Y_0) < \epsilon/(2M)$, and let $Y = p^{-1}(Y_0)$. Then $\eta(Y) = \kappa(Y_0) < \epsilon/M$, and Y is E-invariant so that $\{\varphi_i \restriction Y\}_{i\in\mathbb{N}}$ is an L-graphing of $E \restriction Y$. It follows that

$$C_{\eta\restriction Y}(E \restriction Y) \leqslant \sum_{i\in\mathbb{N}} C_\eta(\{\varphi_i \restriction Y\}) \leqslant M \cdot \eta(Y) + \sum_{i\geqslant M} C_\eta(\{\varphi_i\}) < \epsilon.$$

Claim 2. $E \subseteq E \restriction Y \vee E_{\Phi^{a,A}}$.

Proof. Suppose $(H, f, g)E(H, f', g)$. Since Y_0 is a complete section for E_T there exists $\gamma_1, \ldots, \gamma_k$ and $\epsilon_1, \ldots, \epsilon_k \in \{-1, 1\}$ such that $(\varphi^{s,B}_{\gamma_k})^{\epsilon_k} \circ \cdots (\circ \varphi^{s,B}_{\gamma_1})^{\epsilon_1}((H, g)) \in Y_0$. Let $\gamma = \gamma_k^{\epsilon_k} \cdots \gamma_1^{\epsilon_1}$ and let $(H_0, g_0) =$

$\gamma^s((H,g)) \in Y_0$. It follows that

$$\begin{aligned}\gamma^a(H,f,g) &= (\gamma_k^{\epsilon_k})^a \cdots (\gamma_1^{\epsilon_1})^a(H,f,g) = (\varphi_{\gamma_k}^{a,A})^{\epsilon_k} \circ \cdots \circ (\varphi_{\gamma_1}^{a,A})^{\epsilon_1}(H,f,g)\\ \gamma^a(H,f',g) &= (\gamma_k^{\epsilon_k})^a \cdots (\gamma_1^{\epsilon_1})^a(H,f',g) = (\varphi_{\gamma_k}^{a,A})^{\epsilon_k} \circ \cdots \circ (\varphi_{\gamma_1}^{a,A})^{\epsilon_1}(H,f',g).\end{aligned}$$

This shows that $(H,f,g)E_{\Phi^{a,A}}\gamma^a(H,f,g)$ and $\gamma^a(H,f',g)E_{\Phi^{a,A}}(H,f',g)$. As $\gamma^a(H,f,g) = (H_0, \gamma^{b_H} f, g_0) \in Y$ and $\gamma^a(H,f',g) = (H_0, \gamma^{b_H} f', g_0) \in Y$ we will be done if we can show these two points are E-related. Let $h \in H$ be such that $h^{b_H} f = f'$ and let $k = \gamma h \gamma^{-1}$. Then $k \in \gamma H \gamma^{-1} = H_0$ and

$$k^a(H_0, \gamma^{b_H} f, g_0) = (k\gamma)^a(H,f,g) = (\gamma h)^a(H,f,g) = \gamma^a(H,f',g) = (H_0, \gamma^{b_H} f', g_0)$$

which shows that $(H_0, \gamma^{b_H} f, g_0)E_{(H_0,g_0)}(H_0, \gamma^{b_H} f', g_0)$. □[Claim 2]

We have $C_\eta(E \upharpoonright Y \vee E_{\Phi^{a,A}}) \leqslant 1+\epsilon$. Since we have shown that $E \subseteq E \upharpoonright Y \vee E_{\Phi^{a,A}}$ and that E is an aperiodic normal subequivalence relation of E_a, it follows from [KM05, 24.10] that $C_\eta(E_a) \leqslant C_\eta(E \upharpoonright Y \vee E_{\Phi^{a,A}}) \leqslant 1+\epsilon$. As $\epsilon > 0$ was arbitrary it follows that $C_\eta(E_a) = 1$ and therefore $C(\Gamma) = 1$. □

6.5 Fixed price 1 and shift-minimality

The following lemma will be needed for Theorem 6.36.

Lemma 6.34. *Let θ be an invariant random subgroup of a countable group Γ that concentrates on the infinite amenable subgroups of Γ. Let $\boldsymbol{a} = \Gamma \curvearrowright^a (X,\mu)$ be a free measure preserving action of Γ and let*

$$\boldsymbol{\theta} \times \boldsymbol{a} = \Gamma \curvearrowright^{c \times a} (\mathrm{Sub}_\Gamma \times X, \theta \times \mu)$$

be the product Γ-system. Then $C_{\theta\times\mu}(E_{c\times a}) = 1$.

Remark 6.35. The proof shows that the hypothesis that θ is amenable can be weakened to the hypothesis that θ concentrates on groups of fixed price 1.

Proof. The proof is similar to that of Lemma 6.33. Since $E_{c\times a}$ is aperiodic it suffices to show that $C_{\theta\times\mu}(E_{c\times a}) \leqslant 1$. For each $H \in \mathrm{Sub}_\Gamma$ let $E_{a\upharpoonright H}$ denote the orbit equivalence relation on X generated by $\boldsymbol{a} \upharpoonright H = H \curvearrowright^a (X,\mu)$. Define the subrelation $E \subseteq E_{c\times a}$ on $\mathrm{Sub}_\Gamma \times X$ by $E = \{((H,x),(H,y)) : xE_{a\upharpoonright H}y\}$, i.e.,

$$(H,x)E(L,y) \;\Leftrightarrow\; H = L \text{ and } (\exists h \in H)\,(h^a \cdot x = y).$$

Then E is a normal sub-equivalence relation of $E_{c\times a}$. Since θ concentrates on the infinite subgroups of Γ, E is aperiodic on a $(\theta\times\mu)$-conull set. By [KM05, 24.10] and then [KM05, Proposition 18.4] we therefore have

$$C_{\theta\times\mu}(E_{c\times a}) \leqslant C_{\theta\times\mu}(E) = \int_H C_\mu(E_{a\upharpoonright H})\,d\theta(H) = 1$$

where the last equality follows from [KM05, Corollary 31.2] since θ-almost every H is infinite amenable. □

Theorem 6.36. *Let Γ be a countably infinite group that contains no non-trivial finite normal subgroup. If Γ is not shift-minimal then Γ has fixed price* 1.

Proof. Suppose that Γ is not shift-minimal. By Corollary 3.14 either Γ has a non-trivial normal amenable subgroup N that is necessarily infinite by our hypothesis on Γ, or there is an infinitely generated amenable invariant random subgroup θ of Γ that is weakly contained in $\boldsymbol{s}_\Gamma$. In the first case define $\theta = \delta_N$, so that in either case θ concentrates on the infinite amenable subgroups of Γ, and $\boldsymbol{\theta} \prec \boldsymbol{s}_\Gamma$.

Let (X,μ) denote the underlying measure space of $\boldsymbol{s}_\Gamma$ and consider the product Γ-system

$$\boldsymbol{\theta} \times \boldsymbol{s}_\Gamma = \Gamma \curvearrowright^{c\times s} (\mathrm{Sub}_\Gamma \times X, \theta \times \mu).$$

By Lemma 6.34 we have $C(\boldsymbol{\theta} \times \boldsymbol{s}_\Gamma) = 1$. The action $\boldsymbol{\theta}$ is weakly contained in $\boldsymbol{s}_\Gamma$, so $\boldsymbol{\theta} \times \boldsymbol{s}_\Gamma$ is weakly equivalent to $\boldsymbol{s}_\Gamma$. This implies that Γ has fixed price 1 by (3)⇒(1) of Corollary 6.24. □

Corollary 6.37. *Suppose that Γ does not have fixed price* 1. *Then the following are equivalence*

1. Γ *is shift-minimal.*
2. Γ *contains no non-trivial finite normal subgroups.*
3. AR_Γ *is trivial.*

Proof. (3)⇒(2) is obvious. (2)⇒(1) is immediate from Theorem 6.36 by our assumption that Γ does not have fixed price 1. (1)⇒(3) holds in general with no assumptions on Γ. ☐

Corollary 6.38. *Let Γ be any group that does not have fixed price* 1. *Then* AR_Γ *is finite and* $\Gamma/\mathrm{AR}_\Gamma$ *is shift-minimal.*

Proof. Any group containing an infinite normal amenable subgroup has fixed price 1 [KM05, Proposition 35.2]. Therefore $N = \mathrm{AR}_\Gamma$ is finite. Let $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ be a free measure preserving action of Γ of cost $C_\mu(E_{\boldsymbol{a}}) > 1$. The measure preserving action $\boldsymbol{b}$ of Γ/N on the ergodic components of $\boldsymbol{a} \upharpoonright N$ is free, and since N is finite we have $C(\boldsymbol{b}) \geqslant C(\boldsymbol{a}) > 1$. Thus, Γ/N does not have fixed price 1, and $\mathrm{AR}_{\Gamma/N} = \{e\}$ by Proposition B.1. Corollary 6.37 now shows that Γ/N is shift-minimal. ☐

7 Questions

7.1 General implications

A countable group Γ is called C^*-*simple* if the reduced C^*-algebra of Γ is simple, i.e., $C^*_r(\Gamma)$ has no non-trivial closed two-sided ideals. As observed in the introduction, there is a strong parallel between shift-minimality and C^*-simplicity. The following characterization of C^*-simplicity of a countable group Γ may be found in [Ha07]. Let λ_Γ denote the left-regular representation of Γ on $\ell^2(\Gamma)$.

Proposition 7.1. *Let Γ be a countable group. Then Γ is C^*-simple if and only if $\pi \prec \lambda_\Gamma$ implies $\pi \sim \lambda_\Gamma$ for all nonzero unitary representations π of Γ.*

In this characterization of C^*-simplicity we may actually restrict our attention to irreducible representations of Γ. That is, Γ is C^*-simple if and only if every irreducible unitary representation π of Γ that is weakly contained in λ_Γ is actually weakly equivalent to λ_Γ. See [BH00]. See also [BHV08, Appendix F] and [Di77] for more on weak containment of unitary representations.

Characterization (6) of shift-minimality from Proposition 3.2 also has an analogue for C^*-simplicity. Let $\mathcal{H}$ be an infinite dimensional separable Hilbert space and let $\mathrm{Irr}_\lambda(\Gamma, \mathcal{H})$ denote the Polish space of irreducible representation of Γ on $\mathcal{H}$ that are weakly contained in λ_Γ (see [Di77]). Let $\mathcal{U}(\mathcal{H})$ be the Polish group of all unitary operators on $\mathcal{H}$. Then Γ is C^*-simple if and only if Γ is ICC and the conjugation action of $\mathcal{U}(\mathcal{H})$ on $\mathrm{Irr}_\lambda(\Gamma, \mathcal{H})$ is minimal (i.e., every orbit is dense). See [Ke10, Appendix H.(**C**)].

Consider now the following properties of a countable group Γ:

(UT) Γ has the unique trace property.

(CS) Γ is C^*-simple.

(SM) Γ is shift-minimal.

(UIRS_0) Γ has no non-trivial amenable invariant random subgroup that is weakly contained in $\boldsymbol{s}_\Gamma$.

(UIRS) Γ has no non-trivial amenable invariant random subgroups.

(AR_e) Γ has no non-trivial amenable normal subgroups, i.e., the amenable radical AR_Γ of Γ is trivial.

All of the known implications (besides (SM)$\Leftrightarrow$(UIRS$_0$)) are depicted in Figure 1 in the introduction. It is known that (UT) and (CS) imply (AR$_e$) ([PS79], see also [BH00, Proposition 3]), though it is an open question whether there are any other implications among the properties (UT), (CS), and (AR$_e$) in general [BH00]. The following questions concern some of the remaining implications.

The implication (UT)$\Rightarrow$(SM) was shown in Theorem 5.15. One of the most pressing questions is:

Question 7.2. Does (CS) imply (SM)? That is, are C*-simple groups shift-minimal?

For a positive answer to Question 7.2 it would suffices by Corollary 3.14 to show that if θ is a non-atomic self-normalizing amenable IRS of a countable group Γ that is weakly contained in $\boldsymbol{s}_\Gamma$ then the tracial state on $C^*_r(\Gamma)$ extending φ_θ from the proof of Theorem 5.14 is not faithful.

The implication from (UT) to (UIRS) is quite direct. The converse would mean that a tracial state on $C^*_r(\Gamma)$ different from τ_Γ somehow gives rise to a non-trivial amenable invariant random subgroup of Γ. This is addressed by the following question:

Question 7.3. Does (UIRS) imply (UT)? That is, if Γ does not have any non-trivial amenable invariant random subgroups then does $C^*_r(\Gamma)$ have a unique tracial state?

We know from Theorem 3.16 that (SM) and (UIRS$_0$) are equivalent. The equivalence of (SM) and (UIRS) is open however (clearly though (UIRS)$\Rightarrow$(UIRS$_0$))

Question 7.4. Does (UIRS$_0$) imply (UIRS)?

To obtain a positive answer to Question 7.4 it would be enough to show the following: ($\star$) Every ergodic amenable invariant random subgroup of a countable group Γ that is not almost ascendant is weakly contained in $\boldsymbol{s}_\Gamma$.

Indeed, assume that ($\star$) holds and suppose that Γ does not have (UIRS), i.e., there is an amenable invariant random subgroup θ of Γ other than $\delta_{\langle e \rangle}$. By moving to an ergodic component of θ we may assume without loss of generality that θ is ergodic. If θ is not almost ascendant then ($\star$) implies that $\boldsymbol{\theta}$ is weakly contained in $\boldsymbol{s}_\Gamma$, which shows that Γ does not have (UIRS$_0$). On the other hand, if θ is almost ascendant then, by Corollary B.4, θ concentrates on the subgroups of AR_Γ, and in particular AR_Γ is non-trivial, so $\delta_{\mathrm{AR}_\Gamma}$ witnesses that Γ does not have (UIRS$_0$).

The implication (SM)$\Rightarrow$(AR$_e$) is shown in Proposition 3.15 above. The converse is a tantalizing question:

Question 7.5. Does (AR$_e$) imply (SM)? That is, if Γ has no non-trivial amenable normal subgroup then is every non-trivial m.p. action that is weakly contained in $\boldsymbol{s}_\Gamma$ free?

To obtain a positive answer to Question 7.5 by Corollary 3.14 it would be enough to show that if θ is a non-atomic self-normalizing invariant random subgroup weakly contained in $\boldsymbol{s}_\Gamma$ then θ concentrates on subgroups of the amenable radical of Γ. (Note that θ does indeed concentrate on the amenable subgroups of Γ by NA-ergodicity.)

7.2 Cost and pseudocost

In the infinitely generated setting it appears that pseudocost, rather than cost, may be a more useful way to define an invariant. In addition to the properties exhibited in §6.2, pseudocost enjoys many of the nice properties already known to hold for cost. For instance, pseudocost respects ergodic decomposition, and $\mathrm{PC}(\Gamma) \leqslant \mathrm{PC}(N)$ whenever N is an infinite normal subgroup of Γ. (The proofs are routine: for the first statement one uses the corresponding fact about cost along with basic properties of pseudocost, and the proof of the second is nearly identical to the corresponding proof for cost.)

Question 7.6. Is there an example of a m.p. countable Borel equivalence relation E such that $PC_\mu(E) < C_\mu(E)$?

By Corollary 6.8.(1) the equality $PC_\mu(E) = C_\mu(E)$ holds whenever $C_\mu(E) < \infty$, so the question is whether it is possible to have $PC_\mu(E) < \infty$ and $C_\mu(E) = \infty$. Equivalently: does there exist an increasing sequence $E_0 \subseteq E_1 \subseteq \cdots$, of m.p. countable Borel equivalence relations on (X, μ) with $\mathbf{sup}_n C_\mu(E_n) < \infty$ and $C_\mu(\bigcup_n E_n) = \infty$? If such a sequence $(E_n)_{n\in\mathbb{N}}$ exists then, letting $E = \bigcup_n E_n$, Corollary 6.8.(2) implies that E could not be treeable. In addition, E would provide an example of strict inequality $\beta_1(E)+1 < C_\mu(E)$. This follows from [Ga02, 5.13, 3.23]. Gaboriau has shown that any aperiodic m.p. countable Borel equivalence R satisfies $\beta_1(R) + 1 \leqslant C_\mu(R)$ [Ga02], although it is open whether this inequality can ever be strict. Note that a positive answer to 7.6 would not necessarily provide a counterexample to the fixed price conjecture, even if the equivalence relation E comes from a free action of some group Γ; at this time there is no way to rule out the possibility that such a Γ has fixed cost ∞ while at the same time admitting various free actions with finite pseudocost.

Question 7.7. Suppose that a countable group Γ has some free action $\boldsymbol{a}$ with $C_\mu(\boldsymbol{a}) = \infty$. Does it follow that $C_\mu(\boldsymbol{s}_\Gamma) = \infty$?

By Corollary 6.20, $\boldsymbol{s}_\Gamma$ attains the maximum pseudocost among free actions of Γ. Corollary 6.22 implies that

$$C(\boldsymbol{s}_\Gamma) \geqslant \mathbf{sup}\{C(\boldsymbol{b}) \, : \, \boldsymbol{b} \in \mathrm{FR}(\Gamma, X, \mu) \text{ and either } C(\boldsymbol{b}) < \infty \text{ or } E_b \text{ is treeable}\}.$$

This is not enough to conclude that $\boldsymbol{s}_\Gamma$ always attains the maximum cost among free actions of Γ. A positive answer to Question 7.7 would imply that $\boldsymbol{s}_\Gamma$ always attains this maximum cost.

It would be just as interesting if $\boldsymbol{s}_\Gamma$ could detect whether $C(\Gamma) < \infty$.

Question 7.8. Suppose that a countable group Γ has some free action $\boldsymbol{a}$ with $C_\mu(\boldsymbol{a}) < \infty$. Does it follow that $C_\mu(\boldsymbol{s}_\Gamma) < \infty$?

At this time it appears that one cannot rule out any combination of answers to Questions 7.7 and 7.8. A positive answer to both questions would amount to showing that no group has both free actions of infinite cost and free actions of finite cost – this would essentially affirm a special case of the fixed price conjecture!

7.3 Other questions

It is shown in [T-D12b] that the natural analogue of Question 7.5, where "amenable" is replaced by "finite" and "weakly contained in" is replaced by "is a factor of," has a positive answer:

Theorem 7.9 (Corollary 1.6 of [T-D12b]). *Let Γ be a countable group. If Γ has no non-trivial finite normal subgroups then every non-trivial totally ergodic action of Γ is free.*

In particular, if Γ has no non-trivial finite normal subgroups then every non-trivial factor of $\boldsymbol{s}_\Gamma$ is free.

Here, a measure preserving action of Γ is called *totally ergodic* if all infinite subgroups of Γ act ergodically. Theorem 7.9 motivates the following question concerning strong NA-ergodicity.

Question 7.10. Let $\Gamma \curvearrowright^a (X, \mu)$ be a non-trivial measure preserving action of a countable group Γ. Suppose that for each non-amenable subgroup $\Delta \leqslant \Gamma$ the action $\Delta \curvearrowright^a (X, \mu)$ is strongly ergodic. Does it follow that the stabilizer of almost every point is contained in the amenable radical of Γ?

A positive answer to 7.10 would imply a positive answer to 7.5 by Proposition 3.10.

The following question concerns the converse of Proposition 4.6:

Question 7.11. Suppose Γ is shift-minimal. Is it true that every finite index subgroup of Γ is shift-minimal?

Question 7.11 is equivalent to the question of whether every finite index *normal* subgroup N of a shift-minimal group Γ is shift-minimal. Indeed, suppose the answer is positive for normal subgroups and let K be a finite index subgroup of a shift-minimal group Γ. Then K is ICC, since the ICC property passes to finite index subgroups. Since the group $N = \bigcap_{\gamma \in \Gamma} \gamma K \gamma^{-1}$ is finite index and normal in Γ, it is shift-minimal by our assumption. Proposition 4.6 then implies that K is shift-minimal.

Corollary 4.8 provides a positive answer to Question 7.11 for finite index subgroups which are torsion-free. Theorem 4.7 gives a positive answer for finite index normal subgroups N of Γ for which there is no infinite locally finite invariant random subgroup that is weakly contained in $\boldsymbol{s}_N$. Note that a positive answer to the analogue of Question 7.11 for C^*-simplicity was demonstrated in [BH00] (and likewise for the unique trace property).

The results from §6.2 and §6.5 suggest that the following may have a positive answer:

Question 7.12. If an infinite group Γ has positive first ℓ^2-Betti number then is it true that $C^*_r(\Gamma/AR_\Gamma)$ is simple and has a unique tracial state?

There are already partial results in this direction: Peterson and Thom [PT11] have shown a positive answer under the additional assumptions that Γ is torsion free and that every non-trivial element of $\mathbb{Z}\Gamma$ acts without kernel on $\ell^2\Gamma$.

Finally, we record here a question raised earlier in this paper.

(Question 3.9)**.** Let Γ be a countable group acting by automorphisms on a compact Polish group G and assume the action is tempered. Does it follow that the action is weakly contained in $\boldsymbol{s}_\Gamma$? As a special case, is it true that the action $SL_2(\mathbb{Z}) \curvearrowright (\mathbb{T}^2, \lambda^2)$ is weakly contained in $\boldsymbol{s}_{SL_2(\mathbb{Z})}$?

Appendices

Appendix A Invariant random subgroups as subequivalence relations

This first appendix studies *invariant random partitions* of Γ which are a natural generalization of invariant random subgroups. In §A.1 it is shown that every invariant random partition of Γ comes from a pair $(\boldsymbol{a}, F)$ where $\boldsymbol{a}$ is a free m.p. action of Γ and F is a (Borel) subequivalence relation of E_a. It is shown in §A.2 that for an invariant random subgroup any such pair $(\boldsymbol{a}, F)$ will have the property that F is *normalized* by $\boldsymbol{a}$, i.e., γ^a is in the normalizer of the full group of F for every $\gamma \in \Gamma$.

Many of the ideas here are inspired by (and closely related to) the notion of a *measurable subgroup* developed by Bowen-Nevo [BoNe09] and Bowen [Bo11b]. See also Remark A.14.

A.1 Invariant random partitions

By a *partition* of Γ we mean an equivalence relation on Γ. The set $\mathcal{P}_\Gamma$ of all partitions of Γ is a closed subset of $2^{\Gamma\times\Gamma}$ and Γ acts continuously on $\mathcal{P}_\Gamma$ by left translation $\Gamma \curvearrowright^{\ell} \mathcal{P}_\Gamma$, i.e.,

$$(\alpha, \beta) \in \gamma P \iff (\gamma^{-1}\alpha, \gamma^{-1}\beta) \in P$$

for each $\gamma, \alpha, \beta \in \Gamma$ and $P \in \mathcal{P}_\Gamma$. For $P \in \mathcal{P}_\Gamma$ and $\alpha \in \Gamma$ let $[\alpha]_P = \{\beta : (\alpha, \beta) \in P\}$ denote the P-class of α. Then it is easy to check that $\gamma[\alpha]_P = [\gamma\alpha]_{\gamma P}$ for all $\gamma \in \Gamma$.

Definition A.1. An *invariant random partition* of Γ is a translation-invariant Borel probability measure on $\mathcal{P}_\Gamma$.

Remark A.2. Let IRP_Γ denote the space of all invariant random partitions of Γ. This is a convex set that is compact and metrizable in the weak*-topology. Similarly, let IRS_Γ denote the compact convex set of all invariant random subgroups of Γ. There is a natural embedding $\Phi : \mathrm{Sub}_\Gamma \hookrightarrow \mathcal{P}_\Gamma$ that assigns to each $H \in \mathrm{Sub}_\Gamma$ the partition of Γ determined by the right cosets of H, i.e., $[\delta]_{\Phi(H)} = H\delta$ for $\delta \in \Gamma$. Observe that this embedding is Γ-equivariant between the conjugation action $\Gamma \curvearrowright^c \mathrm{Sub}_\Gamma$ and the translation action $\Gamma \curvearrowright^\ell \mathcal{P}_\Gamma$. We thus obtain an embedding $\Phi_* : \mathrm{IRS}_\Gamma \hookrightarrow \mathrm{IRP}_\Gamma$, $\theta \mapsto \Phi_*\theta$.

Suppose now that $F \subseteq X \times X$ is a measure preserving countable Borel equivalence relation on (X, μ) and $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ is a m.p. action of Γ. Each point $x \in X$ determines a partition $P^a_F(x)$ of Γ given by

$$P^a_F(x) = \{(\alpha, \beta) \in \Gamma : \beta^{-1}xF\alpha^{-1}x\}.$$

Note that $P^a_F(x) = P^a_{F \cap E_a}(x)$ for all $x \in X$, so if we are only concerned with properties of P^a_F then we might as well assume that $F \subseteq E_a$.

Proposition A.3. *The map* $x \mapsto P^a_F(x)$ *is equivariant and therefore* $(P^a_F)_*\mu$ *is an invariant random partition of* Γ*.*

Proof. For any $\gamma \in \Gamma$ and $x \in X$ we have

$$(\alpha, \beta) \in P^a_F(\gamma x) \Leftrightarrow \alpha^{-1}\gamma x F \beta^{-1}\gamma x \Leftrightarrow (\gamma^{-1}\alpha, \gamma^{-1}\beta) \in P^a_F(x) \Leftrightarrow (\alpha, \beta) \in \gamma^\ell \cdot P^a_F(x). \qquad \square$$

Proposition A.3 has a converse in a strong sense: given an invariant random partition ρ of Γ there is a *free* m.p. action $\boldsymbol{b} = \Gamma \curvearrowright^b (Y, \nu)$ of Γ and a subequivalence relation $\mathcal{F}$ of E_b with $(P^b_{\mathcal{F}})_*\nu = \rho$. In fact, $\mathcal{F}$ and b can be chosen independently of ρ, with only ν depending on ρ, as we now show. Let $\boldsymbol{\rho}$ denote the m.p. action $\Gamma \curvearrowright^\ell (\mathcal{P}_\Gamma, \rho)$ and let $\boldsymbol{b} = \boldsymbol{\rho} \times \boldsymbol{s}_\Gamma$ (any free action of Γ will work in place of $\boldsymbol{s}_\Gamma$) so that $(Y, \nu) = (\mathcal{P}_\Gamma \times [0, 1]^\Gamma, \rho \times \lambda^\Gamma)$. Define $\mathcal{F} \subseteq Y \times Y$ by

$$(P, x)\mathcal{F}(Q, y) \;\Leftrightarrow\; \exists\gamma \in \Gamma \;(\gamma^{-1} \in [e]_P \text{ and } (\gamma P, \gamma x) = (Q, y)). \tag{A.1}$$

Theorem A.4. *Let* ρ *be an invariant random partition of* Γ *and write* $\boldsymbol{b} = \Gamma \curvearrowright^b (Y, \nu)$ *for the action* $\boldsymbol{\rho} \times \boldsymbol{s}_\Gamma$*. Let* $\mathcal{F}$ *be given by* (A.1)*. Then* $\mathcal{F}$ *is an equivalence relation contained in the equivalence relation* E_b *generated by the* b*, and* $P^b_{\mathcal{F}}((P, x)) = P$ *for* ν*-almost every* $(P, x) \in Y$*. In particular,* $(P^a_{\mathcal{F}})_*\nu = \rho$*.*

Proof of Theorem A.4. It is clear that $\mathcal{F} \subseteq E_b$. We show that $\mathcal{F}$ is an equivalence relation: It is clear that $\mathcal{F}$ is reflexive. To see $\mathcal{F}$ is symmetric, suppose $(P, x)\mathcal{F}(Q, y)$, as witnessed by $\gamma^{-1} \in [e]_P$ with $\gamma P = Q$ and $\gamma x = y$. Then $\gamma \in [e]_{\gamma P} = [e]_Q$ and $(\gamma^{-1}Q, \gamma^{-1}y) = (P, x)$, so $(Q, y)\mathcal{F}(P, x)$. For transitivity, if $(P, x)\mathcal{F}(Q, y)\mathcal{F}(R, z)$ as witnessed by $\gamma^{-1} \in [e]_P$ with $(\gamma P, \gamma x) = (Q, y)$ and $\delta^{-1} \in [e]_Q$ with $(\delta Q, \delta y) = (R, z)$ then $\gamma^{-1} \in [e]_P$ and $\gamma P = Q$ implies $[e]_Q = [e]_{\gamma P} = \gamma[e]_P$. Therefore $\delta^{-1} \in \gamma[e]_P$, i.e., $(\delta\gamma)^{-1} \in [e]_P$ and $(\delta\gamma P, \delta\gamma x)(\delta Q, \delta y) = (R, z)$.

Fix now $(P, x) \in Y$. We show that $P^b_{\mathcal{F}}((P, x)) = P$. For each $\alpha, \beta \in \Gamma$ we have by definition

$$\begin{aligned}(\alpha, \beta) \in P^a_{\mathcal{F}}((P, x)) \;&\Leftrightarrow (\alpha^{-1}P, \alpha^{-1}x)F(\beta^{-1}P, \beta^{-1}x) \\ &\Leftrightarrow\; \exists\gamma \in \Gamma \left(\gamma^{-1} \in [e]_{\alpha^{-1}P} \text{ and } (\gamma\alpha^{-1}P, \gamma\alpha^{-1}x) = (\beta^{-1}Q, \beta^{-1}x)\right).\end{aligned} \tag{A.2}$$

Therefore, if $(\alpha, \beta) \in P^b_{\mathcal{F}}((P, x))$ as witnessed by some γ as in (A.2) then $\gamma\alpha^{-1}x = \beta^{-1}x$ so freeness of $\boldsymbol{a}$ implies $\gamma = \beta^{-1}\alpha$. Then $\alpha^{-1}\beta = \gamma^{-1} \in [e]_{\alpha^{-1}P}$, i.e., $(\alpha^{-1}\beta, e) \in \alpha^{-1}P$, which is equivalent to $(\beta, \alpha) \in P$. This shows that $P^b_{\mathcal{F}}((P, x)) \subseteq P$. For the reverse inclusion, if $(\alpha, \beta) \in P$ then $\gamma = \beta^{-1}\alpha$ satisfies (A.2) and thus $(\alpha, \beta) \in P^b_{\mathcal{F}}((P, x))$. $\square$

Definition A.5. Let $\boldsymbol{a} = \Gamma \curvearrowright^{a} (X, \mu)$ be a m.p. action of Γ and let F be a subequivalence relation of E_a. If ρ is an invariant random partition of Γ then the pair $(\boldsymbol{a}, F)$ is called a *realization* of ρ if $(P^a_F)_*\mu = \rho$. If θ is an invariant random subgroup of Γ then $(\boldsymbol{a}, F)$ is called a *realization* of θ if it is a realization of $\Phi_*\theta$, where $\Phi_* : \mathrm{IRS}_\Gamma \to \mathrm{IRP}_\Gamma$ is the embedding defined in Remark A.2. A realization $(\boldsymbol{a}, F)$ is called *free* if $\boldsymbol{a}$ is free.

The following is a straightforward consequence of Theorem A.4 and the definitions.

Corollary A.6. *Every invariant random partition admits a free realization.*

The remainder of this subsection works toward a characterization of the set $\Phi_*(\mathrm{IRS}_\Gamma)$. Let K be a metrizable compact space and consider the set $\mathcal{P}_\Gamma \otimes K$ of all pairs (P, f) where $f : P^* \to K$ is a function with $\mathrm{dom}(f) = P^* = \{[\alpha]_P : \alpha \in \Gamma\}$ and taking values in K. The set $\mathcal{P}_\Gamma \otimes K$ has a natural compact metrizable topology coming from its identification with the closed set

$$\widetilde{\mathcal{P}_\Gamma \otimes K} = \{(P, g) \in \mathcal{P}_\Gamma \times K^\Gamma \,:\, g \text{ is constant on each } P\text{-class}\} \subseteq \mathcal{P}_\Gamma \times K^\Gamma$$

via the injection $(P, f) \mapsto (P, \tilde{f})$ where $\tilde{f}(\alpha) = f([\alpha]_P)$ for $\alpha \in \Gamma$. Observe that $\widetilde{\mathcal{P}_\Gamma \otimes K}$ is invariant in $\mathcal{P}_\Gamma \times K^\Gamma$ with respect to the product action $\ell \times s$ of Γ (where s denotes the shift action $\Gamma \curvearrowright^{s} K^\Gamma$), so we obtain a continuous action $\Gamma \curvearrowright^{\ell \otimes s} \mathcal{P}_\Gamma \otimes K$. Explicitly, this action is given by $\gamma \cdot (P, f) = (\gamma P, \gamma^{s_P} f)$ where $\gamma^{s_P} f : (\gamma P)^* \to K$ is the function

$$(\gamma^{s_P} f)([\alpha]_{\gamma P}) = f(\gamma^{-1}[\alpha]_{\gamma P}) = f([\gamma^{-1}\alpha]_P).$$

There is a natural equivalence relation $\mathcal{R} = \mathcal{R}_K$ on $\mathcal{P}_\Gamma \otimes K$ given by

$$(P, f)\mathcal{R}(Q, g) \;\Leftrightarrow\; \exists \gamma \in [e]_P \;(\gamma^{-1}(P, f) = (Q, g)).$$

It is clear that $\mathcal{R}$ is an equivalence relation that is contained in $E_{\ell \otimes s}$.

Lemma A.7. $P \subseteq P^{\ell \otimes s}_{\mathcal{R}}((P, f))$ *for every* $(P, f) \in \mathcal{P}_\Gamma \otimes K$.

Proof. Suppose that $(\alpha, \beta) \in P$. Then $\beta^{-1}\alpha \in [e]_{\beta^{-1}P}$ so for any $f \in K^{P^*}$, from the definition of $\mathcal{R}$ we have

$$(\beta^{-1}P, \beta^{-1}f)\, \mathcal{R}\, (\beta^{-1}\alpha)^{-1}(\beta^{-1}P, \beta^{-1}f) = (\alpha^{-1}P, \alpha^{-1}f),$$

i.e., $\beta^{-1}(P, f)\mathcal{R}\alpha^{-1}(P, f)$. This means that $(\alpha, \beta) \in P^{\ell \otimes s}_{\mathcal{R}}((P, f))$ by definition. □

If ρ is an invariant random partition and μ is a Borel probability measure on K then the measure $\rho \otimes \mu$ on $\mathcal{P}_\Gamma \otimes K$ given by

$$\rho \otimes \mu = \int_P (\delta_P \times \mu^{P^*})\, d\rho$$

is $\ell \otimes s$-invariant.

Theorem A.8. *Let* ρ *be an invariant random partition of* Γ*, let* μ *be any atomless measure on* K*, and let* $\mathcal{R} = \mathcal{R}_K$*. Then the following are equivalent:*

1. $\rho \in \Phi_*(\mathrm{IRS}_\Gamma)$
2. $(\rho \otimes \mu)$*-almost every* $\mathcal{R}$*-class is trivial.*

Proof. (1)⇒(2): Suppose that (1) holds. It follows that $(\rho \otimes \mu)$ concentrates on pairs $(\Phi(H), f) \in \mathcal{P}_\Gamma \otimes K$ with $H \in \mathrm{Sub}_\Gamma$. It therefore suffices to show that the $\mathcal{R}$-class of such a pair $(\Phi(H), f)$ is trivial. If $(\Phi(H), f)\mathcal{R}(Q, g)$ then there is some $\gamma \in [e]_{\Phi(H)} = H$ with $\gamma^{-1}\Phi(H) = Q$ and $\gamma^{-1}f = Q, g$. But $\gamma^{-1}\Phi(H) = \Phi(\gamma^{-1}H\gamma) =$

$\Phi(H)$ (since $\gamma \in H$) so that $Q = \Phi(H)$. In addition, for each $\delta \in \Gamma$ we have $\gamma[\delta]_{\Phi(H)} = \gamma H\delta = H\delta = [\delta]_{\Phi(H)}$ since $\gamma \in H$. Therefore $g([\delta]_{\Phi(H)}) = (\gamma^{-1}f)([\delta]_{\Phi(H)}) = f(\gamma[\delta]_{\Phi(H)}) = f([\delta]_{\Phi(H)})$, showing that $g = f$.

(2)⇒(1): Suppose that (2) holds. Since μ is non-atomic, for each $P \in \mathcal{P}_\Gamma$ the set $\{f \in K^{P^*} : f \text{ is injective}\}$ is μ^{P^*}-conull. This along with (2) implies that there is a Γ-invariant $(\rho\otimes\mu)$-conull set $Y \subseteq \mathcal{P}_\Gamma \otimes K$ on which $\mathcal{R}$ is trivial and such that $f : P^* \to K$ is injective whenever $(P, f) \in Y$. The projection $Y_0 = \{P \in \mathcal{P}_\Gamma : \exists f\ (P, f) \in Y\}$ is then ρ-conull so it suffices to show that $Y_0 \subseteq \Phi(\mathrm{Sub}_\Gamma)$. Fix $P \in Y_0$ and an $f : P^* \to K$ with $(P, f) \in Y$.

Claim 3. *Let $\alpha, \beta \in \Gamma$. Then $(\alpha, \beta) \in P$ if and only if $\beta\alpha^{-1} \in [e]_P$.*

Proof of Claim. Suppose $(\alpha, \beta) \in P$. Lemma A.7 implies $(\alpha, \beta) \in P^a_{\mathcal{R}}(P, f)$ so as the relevant $\mathcal{R}$-classes are trivial this implies $\alpha^{-1}(P, f) = \beta^{-1}(P, f)$ and thus $\alpha\beta^{-1}P = P$ and $\alpha\beta^{-1}f = f$. Then $f([e]_P) = (\alpha\beta^{-1}f)([e]_P) = f([\beta\alpha^{-1}]_P)$ so injectivity of f shows that $[\beta\alpha^{-1}]_P = [e]_P$, i.e., $\beta\alpha^{-1} \in [e]_P$.

Conversely, suppose $\beta\alpha^{-1} \in [e]_P$. Then $(\beta\alpha)^{-1}(P, f)\mathcal{R}(P, f)$ by definition of $\mathcal{R}$, and since the $\mathcal{R}$-classes are trivial this implies $(\beta\alpha)^{-1}(P, f) = (P, f)$ and thus $\beta^{-1}(P, f) = \alpha^{-1}(P, f)$. Therefore $f([\beta]_P) = (\beta^{-1}f)([e]_{\beta^{-1}P}) = (\alpha^{-1}f)([e]_{\alpha^{-1}P}) = f([\alpha]_P)$. Since f is injective we conclude that $[\beta]_P = [\alpha]_P$, i.e., $(\alpha, \beta) \in P$. □[Claim]

It is immediate from the claim that $[e]_P$ is a subgroup of Γ and that P is the partition determined by the right cosets of $[e]_P$, i.e., $P = \Phi([e]_P)$. □

A.2 Normalized subequivalence relations

As in the previous section let $F \subseteq X \times X$ be a m.p. countable Borel equivalence relation on (X, μ) and let $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ be a m.p. action of Γ.

Definition A.9. F is said to be *normalized* by $\boldsymbol{a} = \Gamma \curvearrowright^a (X, \mu)$ if there is a conull set $X_0 \subseteq X$ such that

$$xFy \;\Rightarrow\; \gamma x F \gamma y$$

for all $\gamma \in \Gamma$ and $x, y \in X_0$. Equivalently, F is normalized by $\boldsymbol{a}$ if the image of Γ in $\mathrm{Aut}(X, \mu)$ is contained in the normalizer of the full group of F. A realization $(\boldsymbol{a}, F)$ of an invariant random partition ρ of Γ is called *normal* if F is normalized by $\boldsymbol{a}$.

Note that if F is normalized by $\boldsymbol{a}$ then $F \cap E_a$ is normalized by $\boldsymbol{a}$ and $P^a_{F\cap E_a}(x) = P^a_F(x)$ so it makes sense once again to restrict our attention to the case where $F \subseteq E_a$. Define now

$$\Gamma^a_F(x) = \{\gamma \in \Gamma : \gamma^{-1}xFx\}$$

It follows from the definitions that $\Gamma^a_F(x) = [e]_{P^a_F(x)}$.

Proposition A.10. *Let F be a subequivalence relation of E_a. Then the following are equivalent*

1. *F is normalized by $\boldsymbol{a}$.*
2. *For almost all x, $\Gamma^a_F(x)$ is a subgroup of Γ and $P^a_F(x)$ is the partition of Γ determined by the right cosets of $\Gamma^a_F(x)$, i.e.,*
$$(\alpha, \beta) \in P^a_F(x) \;\Leftrightarrow\; \Gamma^a_F(x)\alpha = \Gamma^a_F(x)\beta.$$
for all $\alpha, \beta \in \Gamma$.
3. *$\Gamma^a_F(\gamma x) = \gamma\Gamma^a_F(x)\gamma^{-1}$ for almost all $x \in X$ and all $\gamma \in \Gamma$.*
4. *The set $[e]_P$ is a subgroup of Γ for $(P^a_F)_*\mu$-almost every $P \in \mathcal{P}_\Gamma$ and the map $P \mapsto [e]_P$ is an isomorphism from $\Gamma \curvearrowright^\ell (\mathcal{P}_\Gamma, (P^a_F)_*\mu)$ to $\Gamma \curvearrowright^c (\mathrm{Sub}_\Gamma, (\Gamma^a_F)_*\mu)$.*

Proof. (1)⇒(2): Suppose (1) holds. By ignoring a null set we may assume without loss of generality that $xFy \Rightarrow \gamma x F \gamma y$ for all $x, y \in X$ and $\gamma \in \Gamma$. We have that $e \in \Gamma^a_F(x)$ for all x. If $\gamma \in \Gamma^a_F(x)$ then $\gamma^{-1}xFx$ so by normality we have $xF\gamma x$ and thus $\gamma^{-1} \in \Gamma^a_F(x)$. If in addition $\delta \in \Gamma^a_F(x)$ then $\delta^{-1}xFxF\gamma x$ so that $\delta^{-1}xF\gamma x$ which by normality implies $\gamma^{-1}\delta^{-1}xFx$, i.e., $\delta\gamma \in \Gamma^a_F(x)$. This shows that $\Gamma^a_F(x)$ is a subgroup. It remains to show that $[\delta]_{P^a_F(x)} = \Gamma^a_F(x)\delta$. We have $\gamma \in [\delta]_{P^a_F(x)}$ if and only if $\delta^{-1}xF\gamma^{-1}x$ which by normality is equivalent to $(\delta\gamma^{-1})xFx$, i.e., $\gamma \in \Gamma^a_F(x)\delta$.

(2)⇒(3): Suppose (2) holds. Then for almost all x and all $\gamma, \delta \in \Gamma$ we have

$$\delta \in \Gamma^a_F(\gamma x) \;\Leftrightarrow\; \delta^{-1}\gamma x F \gamma^a x \;\Leftrightarrow\; \gamma^{-1}\delta^{-1}\gamma x F x \;\Leftrightarrow\; \delta \in \gamma\Gamma^a_F(x)\gamma^{-1}.$$

(3)⇒(1): Suppose that (3) holds. Let $X_0 \subseteq X$ be an E_a-invariant conull set such that $\Gamma^a_F(\gamma x) = \gamma\Gamma^a_F(x)\gamma^{-1}$ for all $x \in X_0$ and $\gamma \in \Gamma$. Then for any $x, y \in F$, if xFy then $xE_a y$ so that $y = \delta x$ for some $\delta \in \Gamma$. This means that $\delta^{-1} \in \Gamma^a_F(x)$ and, so for all $\gamma \in \Gamma$ we have $\gamma\delta^{-1}\gamma^{-1} \in \Gamma^a_F(\gamma x)$ and thus

$$\gamma y = (\gamma\delta^{-1}\gamma^{-1})^{-1}(\gamma x)F\gamma x.$$

This shows that F is normalized by $\boldsymbol{a}$.

(2)+(3)⇒(4): Assume (2) and (3) hold. Then the measure $(P^a_F)_*\mu$ concentrates on $\Phi(\mathrm{Sub}_\Gamma)$. It follows that $P \mapsto [e]_P$ is injective on a $(P^a_F)_*\mu$-conull set. By (3) this map is equivariant on a conull set. Since the composition $x \mapsto P^a_F(x) \mapsto [e]_{P^a_F(x)}$ is the same as $x \mapsto \Gamma^a_F(x)$ this map is measure preserving.

Finally, the implication (4)⇒(3) is clear. □

The following corollary is immediate.

Corollary A.11. *If* F *is normalized by* $\boldsymbol{a}$ *then* $(\Gamma^a_F)_*\mu$ *is an invariant random subgroup of* Γ.

Theorem A.4 also implies a converse to Corollary A.11. Let θ be an invariant random subgroup of Γ and let $\rho = \Phi_*\theta$. Let $\boldsymbol{b}$ and $\mathcal{F}$ be defined as in Theorem A.4. Let $\boldsymbol{a} = \boldsymbol{\theta} \times \boldsymbol{s}_\Gamma$ so that $(X, \mu) = (\mathrm{Sub}_\Gamma \times [0,1]^\Gamma, \theta \times \lambda)$. Then the map $\Psi : (H, x) \mapsto (\Phi(H), x)$ is an isomorphism of $\boldsymbol{a}$ with $\boldsymbol{b}$. Letting $\mathcal{F}_0 = (\Psi \times \Psi)^{-1}(\mathcal{F})$, we have that

$$(H, x)\mathcal{F}_0(L, y) \;\Leftrightarrow\; H = L \text{ and } (\exists h \in H)(h^a x = y). \tag{A.3}$$

Corollary A.12. $\mathcal{F}_0$ *is a subequivalence relation of* E_a *on* X *which is normalized by* $\boldsymbol{a}$ *and satisfies* $\Gamma^a_{\mathcal{F}_0}(H, x) = H$ *for* $\theta \times \mu$*-almost-every* $(H, x) \in X$. *Thus* $(P^a_{\mathcal{F}_0})_*\mu = \Phi_*\theta$. *It follows that every invariant random subgroup of* Γ *admits a normal, free realization.*

Proof. All that needs to be checked is that $\mathcal{F}_0$ is normalized by $\boldsymbol{\theta} \times \boldsymbol{a}$. If $(H, x)\mathcal{F}_0(L, y)$ then $H = L$ and $h^a x = y$ for some $h \in H$. Then for any $\gamma \in \Gamma$ we must show that $\gamma \cdot (H, x)\,\mathcal{F}_0\,\gamma \cdot (H, h^a x)$. Now, $\gamma \cdot (H, x) = (\gamma H \gamma^{-1}, \gamma^a x)$, so as $\gamma h \gamma^{-1} \in \gamma H \gamma^{-1}$ the definition (A.3) of $\mathcal{F}_0$ shows that

$$(\gamma H\gamma^{-1}, \gamma^a x)\,\mathcal{F}_0\,\gamma h \gamma^{-1} \cdot (\gamma H \gamma^{-1}, \gamma^a x) = \gamma \cdot (H, h^a x)$$

□

Remark A.13. In Corollary A.12, if θ concentrates on the amenable subgroups of Γ then $\mathcal{F}_0$ will always be an amenable equivalence relation. For other properties of θ, a judicious choice of free action $\boldsymbol{d}$ in place of $\boldsymbol{s}_\Gamma$ in the definition of $\boldsymbol{a}$ may ensure that properties of θ are reflected by the equivalence relation F. For example, if θ concentrates on subgroups of cost r then the proof of Theorem 6.31 above shows that $\boldsymbol{d}$ can be chosen so that the corresponding equivalence relation $\mathcal{F}_0$ has cost r. Similarly, if θ concentrates on treeable subgroups then $\mathcal{F}_0$ can be made a treeable equivalence relation.

Remark A.14. Following [BoNe09, §2.2] let $2^\Gamma_e = \{L \in 2^\Gamma : e \in L\}$ and define the equivalence relation $\mathcal{R}_e \subseteq 2^\Gamma_e \times 2^\Gamma_e$ by

$$(L, K) \in \mathcal{R}_e \Leftrightarrow \exists \gamma \in L \;\; \gamma^{-1}L = K.$$

Then any $\mathcal{R}_e$-invariant Borel probability measure on 2^Γ_e is called a a *measurable subgroup* of Γ (see [BoNe09] and [Bo11b]). If ρ is any invariant random partition of Γ then the image of ρ under $P \mapsto [e]_P$ is a measurable subgroup of Γ. I do not know whether every measurable subgroup of Γ comes from an invariant random partition in this way.

Creutz and Peterson [CP12] define the *subgroup* partial order on $(\mathrm{IRS}_\Gamma, \leqslant)$ as follows: Let $\theta_1, \theta_2 \in \mathrm{IRS}_\Gamma$. Then θ_1 is called a *subgroup* of θ_2 (written $\theta_1 \leqslant \theta_2$) if there exists a joining of θ_1 and θ_2 that concentrates on the set $\{(H, L) \in \mathrm{Sub}_\Gamma : H \leqslant L\}$. It is shown in [CP12] that this is a partial order on IRS_Γ. The same idea can be used to define a notion of refinement for invariant random partitions.

For partitions $P, Q \in \mathcal{P}_\Gamma$, P is said to *refine* Q, written $P \leqslant Q$, if P is a subset of Q. Equivalently $P \leqslant Q$ means that $[\alpha]_P \subseteq [\alpha]_Q$ for every $\alpha \in \Gamma$. If ρ_1 and ρ_2 are invariant random partitions of Γ then ρ_1 *refines* ρ_2, written $\rho_1 \leqslant \rho_2$, if there exists a joining of ρ_1 and ρ_2 that concentrates on the set $\{(P, Q) \in \mathcal{P}_\Gamma \times \mathcal{P}_\Gamma : P \leqslant Q\}$. It is clear that the restriction of the refinement relation on $\mathcal{P}_\Gamma$ (respectively, IRP_Γ) to Sub_Γ (respectively, IRS_Γ) is the subgroup relation.

The point of view developed in this section can be used to give a characterization of the partial orders $(\mathrm{IRS}_\Gamma, \leqslant)$ and $(\mathrm{IRP}_\Gamma, \leqslant)$ in terms of subequivalence relations of free actions of Γ.

Theorem A.15. *Let $\rho_1, \rho_2 \in \mathrm{IRP}_\Gamma$. Then the following are equivalent*

(1) $\rho_1 \leqslant \rho_2$

(2) There exists a free m.p. action $\Gamma \curvearrowright^a (X, \mu)$ of Γ and equivalence relations $F_1 \subseteq F_2 \subseteq E_a$ with $(P^a_{F_1})_*\mu = \rho_1$ and $(P^a_{F_2})_*\mu = \rho_2$.

If $\theta_1, \theta_2 \in \mathrm{IRS}_\Gamma$ then then following are equivalent

(1') $\theta_1 \leqslant \theta_2$.

(2') There exists a free m.p. action $\Gamma \curvearrowright^a (X, \mu)$ of Γ and normalized equivalence relations $F_1 \subseteq F_2 \subseteq E_a$ with $(\Gamma^a_{F_1})_*\mu = \theta_1$ and $(\Gamma^a_{F_2})_*\mu = \theta_2$.

Proof. Suppose (2) holds and let $P^a_{F_1} \times P^a_{F_2} : X \to \mathcal{P}_\Gamma \times \mathcal{P}_\Gamma$ be the map $x \mapsto (P^a_{F_1}(x), P^a_{F_2}(x))$. Then $(P^a_{F_1} \times P^a_{F_2})_*\mu$ is a joining of ρ_1 and ρ_2 with the desired property.

Assume that (1) holds and let ν be a joining of ρ_1 and ρ_2 witnessing that $\rho_1 \leqslant \rho_2$. Let $X = \mathcal{P}_\Gamma \times \mathcal{P}_\Gamma \times [0, 1]^\Gamma$, let $\mu = \nu \times \lambda^\Gamma$, and let $a = \ell \times \ell \times s$. Then we define the equivalence relations F_1 and F_2 on X by

$$(P_1, P_2, x)F_1(Q_1, Q_2, y) \Leftrightarrow \exists \gamma \in \Gamma(\gamma^{-1} \in [e]_{P_1} \text{ and } \gamma^a \cdot (P_1, P_2, x) = (Q_1, Q_2, y))$$
$$(P_1, P_2, x)F_2(Q_1, Q_2, y) \Leftrightarrow \exists \gamma \in \Gamma(\gamma^{-1} \in [e]_{P_2} \text{ and } \gamma^a \cdot (P_1, P_2, x) = (Q_1, Q_2, y)).$$

Then as in the proof of Theorem A.4, F_1 and F_2 are equivalence relations that are contained in E_a and $(\boldsymbol{a}, F_i)$ is a realization of F_i for each $i \in \{1, 2\}$. The defining property of ν also ensures that $F_1 \subseteq F_2$.

The equivalence of (1') and (2') then follows from the equivalence of (1) and (2) along with Proposition A.10. ☐

Finally, we note the following (observed by Vershik [Ve11] in the case of invariant random subgroups), which is a consequence of [IKT09, §1].

Theorem A.16. *Let ρ be an invariant random partition of Γ. Then the function*

$$\varphi_\rho(\gamma) = \rho(\{P : \gamma \in [e]_P\})$$

is a positive definite function on Γ.

Proof. By Corollary A.6 there is a free m.p. action $\boldsymbol{b} = \Gamma \curvearrowright^b (Y, \nu)$ of Γ and a subequivalence relation F of E_b such that $(P^b_F)_*\nu = \rho$. Thus

$$\varphi_\rho(\gamma) = \nu(\{y : \gamma^{-1}yFy\}).$$

This is a positive definite function by [IKT09]. ☐

Appendix B The amenable radical of a countable group

Every countable discrete group Γ contains a largest normal amenable subgroup called the *amenable radical* of Γ (see, e.g., [Zi84, 4.1.12]). We write AR_Γ for the amenable radical of Γ. We present in this appendix some facts concerning AR_Γ for countable Γ.

B.1 Basic properties of AR_Γ

Proposition B.1. *Let Γ be a countable group.*

(1) AR_Γ *is an amenable characteristic subgroup of Γ which contains every normal amenable subgroup of Γ.*

(2) Suppose $\varphi : \Gamma \to \Delta$ is a group homomorphism and that ker(φ) is amenable. Then $\varphi(\mathrm{AR}_\Gamma) = \mathrm{AR}_{\varphi(\Gamma)}$. In particular, the amenable radical of the quotient group $\Gamma/\mathrm{AR}_\Gamma$ is trivial.

(3) If H is normal in Γ then AR_H is a normal subgroup of AR_Γ with $\mathrm{AR}_H = \mathrm{AR}_\Gamma \cap H$.

(4) If H is finite index in Γ then AR_H is a finite index subgroup of AR_Γ with $\mathrm{AR}_H = \mathrm{AR}_\Gamma \cap H$.

Proof. For (1) see [Zi84]. For (2), let $N = \ker(\varphi)$. It is clear that $\varphi(\mathrm{AR}_\Gamma)$ is a normal amenable subgroup of $\varphi(\Gamma)$, so that $\varphi(\mathrm{AR}_\Gamma) \leqslant \mathrm{AR}_{\varphi(\Gamma)}$ by (1). The group $K = \varphi^{-1}(\mathrm{AR}_{\varphi(\Gamma)})$ is normal in Γ and K is amenable since both N and $K/N \cong \mathrm{AR}_{\varphi(\Gamma)}$ are amenable. Hence $K \leqslant \mathrm{AR}_\Gamma$ and so $\mathrm{AR}_{\varphi(\Gamma)} \leqslant \varphi(K) \leqslant \varphi(\mathrm{AR}_\Gamma)$.

We now prove (3). Suppose that H is normal in Γ. It is clear that $\mathrm{AR}_\Gamma \cap H$ is normal in AR_Γ, so it suffices to show that $\mathrm{AR}_\Gamma \cap H = \mathrm{AR}_H$. Conjugation by any element of Γ is an automorphism of H, so fixes (setwise) the characteristic subgroup AR_H. This shows that AR_H is normal in Γ, and since it is amenable it must be contained in AR_Γ. Thus $\mathrm{AR}_H \leqslant \mathrm{AR}_\Gamma \cap H$. In addition, $\mathrm{AR}_\Gamma \cap H$ is a normal amenable subgroup of H, so $\mathrm{AR}_\Gamma \cap H \leqslant \mathrm{AR}_H$. This proves (3).

We need the following Lemma for (4):

Lemma B.2. *Suppose that K is an amenable subgroup of Γ whose normalizer $N_\Gamma(K)$ is finite index in Γ. Then $K \leqslant \mathrm{AR}_\Gamma$.*

Proof of Lemma B.2. Suppose first that K is finite. $N_\Gamma(K)$ being finite index means K has only finitely many conjugates in Γ, so as K itself is finite this implies that every element of K has a finite conjugacy class in Γ. Thus, $K \subseteq \mathrm{FC}_\Gamma \subseteq \mathrm{AR}_\Gamma$, where FC_Γ is the amenable characteristic subgroup of Γ consisting of all elements of Γ with finite conjugacy classes (see e.g., [Ha07, Appendix J]).

Suppose now that K is infinite. The normal core $N = \bigcap_{\gamma \in \Gamma} \gamma N_\Gamma(K)\gamma^{-1}$ of $N_\Gamma(K)$ in Γ is a normal finite index subgroup of Γ. Thus, letting $H = K \cap N$, we have $[K : H] = [KN : N] \leqslant [\Gamma : N] < \infty$, and so H is finite index in K. It is clear that H is normal in N, and H is an amenable group since it is a subgroup of K. Thus $H \leqslant \mathrm{AR}_N$. In addition, AR_N is normal in Γ since AR_N is characteristic in N and N is normal in Γ. Therefore

$$H \leqslant \mathrm{AR}_N \leqslant \mathrm{AR}_\Gamma.$$

Now, H is finite index in K, and $H \leqslant \mathrm{AR}_\Gamma$, so the image $p(K)$ of K in $\Gamma/\mathrm{AR}_\Gamma$ under the quotient map p is a finite subgroup of Γ. So if $p(K)$ were non-trivial then $\Gamma/\mathrm{AR}_\Gamma$ would have non-trivial amenable radical, contrary to part (2). □[Lemma B.2]

We can now show (4). If H is finite index in Γ, then AR_H is an amenable subgroup of Γ whose normalizer $N_\Gamma(\mathrm{AR}_H)$ contains H. Therefore $N_\Gamma(\mathrm{AR}_H)$ is finite index in Γ, so $\mathrm{AR}_H \leqslant \mathrm{AR}_\Gamma$ by Lemma B.2, and thus $\mathrm{AR}_H \leqslant \mathrm{AR}_\Gamma \cap H$. The group AR_Γ is normal in Γ, so $\mathrm{AR}_\Gamma \cap H$ is normal in H and since it is an amenable group we have the other inclusion $\mathrm{AR}_\Gamma \cap H \leqslant \mathrm{AR}_H$. □

Lemma B.3. *Let* Γ *be a countable group and let* $\{H_\alpha\}_{\alpha \leqslant \lambda}$ *be an almost ascendant series in* Γ *(Definition 4.12). Then* $\{AR_{H_\alpha}\}_{\alpha \leqslant \lambda}$ *is an almost ascendant series in* AR_Γ*. The same holds if we replace "almost ascendant" by "ascendant."*

Proof. We show by transfinite induction on ordinals α (with $\alpha \leqslant \lambda$) that $\{AR_{H_\beta}\}_{\beta \leqslant \alpha}$ is an almost ascendant series in AR_{H_α}. If $\alpha = \beta + 1$ is a successor ordinal then by hypothesis H_β is either normal or finite index in $H_{\beta+1}$. Proposition B.1 then implies that AR_{H_β} is either normal or finite index in $AR_{H_{\beta+1}}$.

Suppose now that α is a limit ordinal and let $K = \bigcup_{\beta<\alpha} AR_{H_\beta}$. We must show that $AR_{H_\alpha} = K$. By the induction hypothesis the groups AR_{H_β}, $\beta < \alpha$, are increasing with β, so K is amenable, being an increasing union of amenable groups. Additionally, K is normal in H_α as we now show. For each $h \in H_\alpha$ there is some $\beta_0 < \alpha$ such that $h \in H_{\beta_0}$. Therefore $h \in H_\beta$ for all $\beta_0 \leqslant \beta < \alpha$. Thus h normalizes AR_{H_β} for all $\beta_0 \leqslant \beta < \alpha$, and since the AR_{H_β} are increasing we have

$$hKh^{-1} = \bigcup_{\beta_0 \leqslant \beta < \alpha} hAR_{H_\beta}h^{-1} = \bigcup_{\beta_0 \leqslant \beta < \alpha} AR_{H_\beta} = K.$$

It follows that $K \leqslant AR_{H_\alpha}$. We have the equality $K = AR_{H_\alpha}$ since $AR_{H_\alpha} = \bigcup_{\beta<\alpha}(AR_{H_\alpha} \cap H_\beta) \leqslant \bigcup_{\beta<\alpha} AR_{H_\beta} = K$. □

Corollary B.4. *Let* Γ *be a countable group and let* H *be an almost ascendant subgroup of* Γ*. Then*

$$AR_H = AR_\Gamma \cap H,$$

In particular, AR_H *is contained in* AR_Γ*, and* AR_Γ *contains every almost ascendant amenable subgroup of* Γ*.*

Proof. The containment $AR_H \leqslant AR_\Gamma \cap H$ is immediate from Lemma B.3. We have equality since $AR_\Gamma \cap H$ is an amenable normal subgroup of H. □

Corollary B.5. *Let* Γ *be a countable group and let* $\gamma \in \Gamma$*. If the centralizer* $C_\Gamma(\gamma)$ *of* γ *is almost ascendant in* Γ *then* $\gamma \in AR_\Gamma$*. Thus, if* AR_Γ *is trivial then the centralizer of any non-trivial element of* Γ *is not almost ascendant.*

Proof. The group $\langle\gamma\rangle$ is a normal amenable subgroup of $C_\Gamma(\gamma)$, so if $C_\Gamma(\gamma)$ is almost ascendent then $\langle\gamma\rangle \leqslant AR_{C_\Gamma(\gamma)} \leqslant AR_\Gamma$ by B.4. □

B.2 Groups with trivial amenable radical

Lemma B.6. *Let* N *be a normal subgroup of* Γ*. Then* AR_Γ *is trivial if and only if both* AR_N *and* $AR_{C_\Gamma(N)}$ *are trivial.*

Proof. Since N is normal in Γ, $C_\Gamma(N)$ is normal in Γ as well. Thus, if AR_Γ is trivial it follows from Proposition B.1 that both AR_N and $AR_{C_\Gamma(N)}$ are trivial.

Suppose now that AR_N and $AR_{C_\Gamma(N)}$ are trivial. We have

$$AR_\Gamma \cap N = AR_N = \{e\}$$

and thus AR_Γ and N must commute, being normal subgroups of Γ with trivial intersection. This means that $AR_\Gamma \leqslant C_\Gamma(N)$ and so

$$AR_\Gamma = AR_\Gamma \cap C_\Gamma(N) = AR_{C_\Gamma(N)} = \{e\}.$$ □

Lemma B.7. *Suppose* $\{H_\alpha\}_{\alpha \leqslant \lambda}$ *is an ascendant series of length* λ *and suppose* $\Gamma = H_\lambda$ *has trivial amenable radical. Then* $AR_{C_\Gamma(H_\alpha)} = \{e\}$ *for all* $\alpha \leqslant \lambda$*.*

Proof. We proceed by transfinite induction on λ. By Corollary B.4 we know that $\mathrm{AR}_{H_\alpha} = \{e\}$ for all $\alpha \leqslant \lambda$.

Limit stages: Suppose first that λ is a limit ordinal. Fix $\alpha \leqslant \lambda$ and let $H = H_\alpha$. By intersecting each term of the ascendant series $\{H_\beta\}_{\beta \leqslant \lambda}$ with $C_\Gamma(H)$ we obtain the series $\{C_{H_\beta}(H)\}_{\beta \leqslant \lambda}$ which is ascendant in $C_\Gamma(H)$. Lemma B.3 implies that $\{\mathrm{AR}_{C_{H_\beta}(H)}\}_{\beta \leqslant \lambda}$ is an ascendant series in $\mathrm{AR}_{C_\Gamma(H)}$ and so

$$\mathrm{AR}_{C_\Gamma(H)} = \bigcup_{\alpha \leqslant \beta < \lambda} \mathrm{AR}_{C_{H_\beta}(H)} \tag{B.1}$$

where the union is increasing. For each β with $\alpha \leqslant \beta < \lambda$ the series $\{H_\xi\}_{\xi \leqslant \beta}$ has length strictly less than λ, so by the induction hypothesis we have

$$\mathrm{AR}_{C_{H_\beta}(H)} = \{e\}.$$

Since this holds for each β with $\alpha \leqslant \beta < \lambda$, equation (B.1) shows that $\mathrm{AR}_{C_\Gamma(H)} = \{e\}$ as was to be shown.

Successor stages: Suppose now that $\lambda = \mu + 1$ is a successor ordinal. Fix for the moment some $\alpha < \lambda$ and let $H = H_\alpha$. Applying the induction hypothesis to the ascendant series $\{H_\beta\}_{\beta \leqslant \mu}$ in H_μ we obtain that $\mathrm{AR}_{C_{H_\mu}(H)} = \{e\}$. Since H_μ is normal in Γ, $C_{H_\mu}(H)$ is normal in $C_\Gamma(H)$, so it follows from Proposition B.1.(3) that

$$\mathrm{AR}_{C_\Gamma(H)} \cap H_\mu = \mathrm{AR}_{C_\Gamma(H)} \cap C_{H_\mu}(H) = \mathrm{AR}_{C_{H_\mu}(H)} = \{e\}. \tag{B.2}$$

Since α was an arbitrary ordinal satisfying $\alpha < \lambda$, (B.2) holds for all $\alpha < \lambda$. We use this to show the following.

Claim 4. *Let ξ and β be ordinals with $\xi \leqslant \beta < \lambda$. Then*

$$\mathrm{AR}_{C_\Gamma(H_\xi)} \leqslant \mathrm{AR}_{C_\Gamma(H_\beta)}$$

Proof of Claim 4. We show by transfinite induction on $\beta < \lambda$ that $\{\mathrm{AR}_{C_\Gamma(H_\xi)}\}_{\xi \leqslant \beta}$ is increasing in ξ. If $\beta = 0$ this is trivial. If $\beta = \alpha + 1$ is a successor ordinal then the induction hypothesis tells us that $\{\mathrm{AR}_{C_\Gamma(H_\xi)}\}_{\xi \leqslant \alpha}$ is increasing with ξ and we must show that $\mathrm{AR}_{C_\Gamma(H_\alpha)} \leqslant \mathrm{AR}_{C_\Gamma(H_{\alpha+1})}$.

Since H_α is normal in $H_{\alpha+1}$, Proposition B.1.(2) shows that $H_{\alpha+1}$ normalizes $\mathrm{AR}_{C_\Gamma(H_\alpha)}$. Thus, for $\delta \in H_{\alpha+1}$ and $\gamma \in \mathrm{AR}_{C_\Gamma(H_\alpha)}$ we have

$$\begin{aligned} &(\delta\gamma\delta^{-1})\gamma^{-1} \in \mathrm{AR}_{C_\Gamma(H_\alpha)} \\ &\delta(\gamma\delta^{-1}\gamma^{-1}) \in H_\mu(\gamma H_\mu \gamma^{-1}) = H_\mu \\ \text{so that} \qquad &\delta\gamma\delta^{-1}\gamma^{-1} \in \mathrm{AR}_{C_\Gamma(H_\alpha)} \cap H_\mu = \{e\} \end{aligned}$$

by (B.2) (we use in the second line that $H_{\alpha+1} \leqslant H_\mu$ and $H_\mu \lhd \Gamma$). This shows that the groups $\mathrm{AR}_{C_\Gamma(H_\alpha)}$ and $H_{\alpha+1}$ commute, and so $\mathrm{AR}_{C_\Gamma(H_\alpha)}$ is a subgroup of $C_\Gamma(H_{\alpha+1})$. As $C_\Gamma(H_{\alpha+1})$ is contained in $C_\Gamma(H_\alpha)$ we conclude that $\mathrm{AR}_{C_\Gamma(H_\alpha)}$ is normal in $C_\Gamma(H_{\alpha+1})$ and therefore $\mathrm{AR}_{C_\Gamma(H_\alpha)} \leqslant \mathrm{AR}_{C_\Gamma(H_{\alpha+1})}$.

Now suppose β is a limit ordinal. The induction hypothesis tells us that $\{\mathrm{AR}_{C_\Gamma(H_\xi)}\}_{\xi < \beta}$ is increasing with $\xi < \beta$ and we must show that $\mathrm{AR}_{C_\Gamma(H_\xi)} \leqslant \mathrm{AR}_{C_\Gamma(H_\beta)}$ for all $\xi < \beta$. Fix $\xi < \beta$. For each α with $\xi \leqslant \alpha < \beta$ we have that $\mathrm{AR}_{C_\Gamma(H_\xi)} \leqslant \mathrm{AR}_{C_\Gamma(H_\alpha)} \leqslant C_\Gamma(H_\alpha)$. Intersecting this over all such α shows

$$\mathrm{AR}_{C_\Gamma(H_\xi)} \leqslant \bigcap_{\xi \leqslant \alpha < \beta} C_\Gamma(H_\alpha) = C_\Gamma\Big(\bigcup_{\xi \leqslant \alpha < \beta} H_\alpha\Big) = C_\Gamma(H_\beta).$$

Since $C_\Gamma(H_\beta) \leqslant C_\Gamma(H_\xi)$ we actually have $\mathrm{AR}_{C_\Gamma(H_\xi)} \lhd C_\Gamma(H_\beta)$ and so $\mathrm{AR}_{C_\Gamma(H_\xi)} \leqslant \mathrm{AR}_{C_\Gamma(H_\beta)}$, which finishes the proof of the claim. □[Claim 4]

Given now any $\alpha < \lambda$ we have shown that $\mathrm{AR}_{C_\Gamma(H_\alpha)} \leqslant \mathrm{AR}_{C_\Gamma(H_\mu)}$. But H_μ is normal in Γ and $\mathrm{AR}_\Gamma = \{e\}$, so Lemma B.6 shows that $\mathrm{AR}_{C_\Gamma(H_\mu)} = \{e\}$ and therefore $\mathrm{AR}_{C_\Gamma(H_\alpha)} = \{e\}$ as was to be shown. □[Lemma B.7]

Lemma B.8. *Let* $\{H_\alpha\}_{\alpha\leqslant\lambda}$ *be an ascendant series of length* λ *with* $H_0 = H$ *and* $H_\lambda = \Gamma$. *Suppose that* $AR_{C_\Gamma(H)} = AR_H = \{e\}$. *Then* $AR_\Gamma = \{e\}$.

Proof. We proceed by transfinite induction on the length λ of the series.

Limit stages: Suppose first that λ is a limit ordinal. By intersecting each group in the series $\{H_\alpha\}_{\alpha\leqslant\lambda}$ with $C_\Gamma(H)$ we obtain the series $\{C_{H_\alpha}(H)\}_{\alpha\leqslant\lambda}$, which is ascendant in $C_\Gamma(H)$. Applying Lemma B.3 to the series $\{C_{H_\alpha}(H)\}_{\alpha\leqslant\lambda}$ we obtain

$$\bigcup_{\alpha<\lambda} AR_{C_{H_\alpha}(H)} = AR_{C_\Gamma(H)}.$$

Since $AR_{C_\Gamma(H)} = \{e\}$ we conclude that $AR_{C_{H_\alpha}(H)} = \{e\}$ for all $\alpha < \lambda$. In addition we have $AR_H = \{e\}$ so it follows from the induction hypothesis (applied to each series $\{H_\xi\}_{\xi<\alpha}$ for $\alpha < \lambda$) that $AR_{H_\alpha} = \{e\}$ for all α. Another application of Lemma B.3 now shows that $AR_\Gamma = \bigcup_{\alpha<\lambda} AR_{H_\alpha} = \{e\}$.

Successor stages: Now assume that $\lambda = \mu + 1$ is a successor ordinal. Since H_μ is normal in $H_{\mu+1} = \Gamma$ we have $C_{H_\mu}(H) \lhd C_\Gamma(H)$. It follows that $AR_{C_{H_\mu}(H)} \leqslant AR_{C_\Gamma(H)} = \{e\}$ and so

$$AR_{C_{H_\mu}(H)} = \{e\}.$$

By assumption $AR_H = \{e\}$ so the induction hypothesis applied to $\{H_\alpha\}_{\alpha\leqslant\mu}$ implies that

$$AR_{H_\mu} = \{e\}. \tag{B.3}$$

Since H_μ is normal in Γ, $C_\Gamma(H_\mu)$ is normal in Γ as well. In addition, $C_\Gamma(H_\mu)$ is contained in $C_\Gamma(H)$, so in fact $C_\Gamma(H_\mu) \lhd C_\Gamma(H)$. It follows that

$$AR_{C_\Gamma(H_\mu)} \leqslant AR_{C_\Gamma(H)} = \{e\}. \tag{B.4}$$

We see from (B.3) and (B.4) that the normal subgroup H_μ of Γ satisfies the hypotheses of Lemma B.6 and so $AR_\Gamma = \{e\}$. This completes the induction. ☐

Theorem B.9. *Let* H *be an ascendant subgroup of a countable group* Γ. *Then* $AR_\Gamma = \{e\}$ *if and only if* $AR_H = \{e\}$ *and* $AR_{C_\Gamma(H)} = \{e\}$.

References

[AGV12] M. Abért, Y. Glasner, and B. Virag, *Kesten's Theorem for Invariant Random Subgroups*, arXiv:1201.3399v1, 2012.

[AW11] M. Abért and B. Weiss, *Bernoulli actions are weakly contained in any free action.* arXiv:1103.1063v1, 2011.

[AM07] G. Arzhantseva and A. Minasyan, *Relatively hyperbolic groups are* C^**-simple.* J. Funct. Anal. 243 (2007), no. 1, 345-351.

[Be91] E. Bèdos, *Discrete groups and simple* C^**-algebras.* Math. Proc. Cambridge Philos. Soc. 109 (1991), no. 3, 521-537.

[BCH94] B. Bekka, M. Cowling and P. de la Harpe, *Some groups whose reduced* C^**-algebra is simple.* Inst. Hautes Études Sci. Publ. Math. No. 80 (1994), 117-134.

[BH00] B. Bekka and P. de la Harpe, *Groups with Simple Reduced* C^**-algebras.* Expo. Math. 18 (2000), no. 3, 215-230.

[BHV08] B. Bekka, P. de la Harpe, A. Valette, *Kazhdan's property (T)*, New Mathematical Monographs, 11. Cambridge University Press, Cambridge, 2008.

[Ber96] V. Bergelson, *The multifarious Poincaré recurrence theorem*, in "Descriptive set theory and dynamical systems," (Marseille-Luminy, 1996), 31-57, London Math. Soc. Lecture Note Ser., 277, Cambridge Univ. Press, Cambridge, 2000.

[BG05] N. Bergeron and D. Gaboriau, *Asymptotique des nombres de Betti, invariants l2 et laminations* Comment. Math. Helv. 79 (2004), no. 2, 362-395.

[BN88] F. Boca and V. Nitica, *Combinatorial properties of groups and simple* C^**-algebras*, J. Operator Theory 20 (1988) 183-196.

[Bo11a] L. Bowen, *Entropy for expansive algebraic actions of residually finite groups*, Ergodic Theory Dynam. Systems 31 (2011), no. 3, 703-718.

[Bo11b] L. Bowen, *Every countably infinite group is almost Ornstein*, Dynamical systems and group actions, 67-78, Contemp. Math., 567, Amer. Math. Soc., Providence, RI, 2012.

[BoNe09] L. Bowen and A. Nevo, *Geometric covering arguments and ergodic theorems for free groups*, LŠEnseignement Mathématique (2009).

[CP12] D. Creutz and J. Peterson, *Stabilizers of Ergodic Actions of Lattices and Commensurators*, preprint (2012).

[CKT-D12] C. Conley, A. Kechris, and R. Tucker-Drob, *Ultraproducts of measure preserving actions and graph combinatorics*, Ergodic Theory and Dynam. Systems, 2012.

[Di77] J. Dixmier, C^**-algebras*, North-Holland Mathematical Library, Vol. 15. North-Holland Publishing Co., Amsterdam-New York-Oxford, 1977.

[FW04] M. Foreman and B. Weiss, *An anti-classification theorem for ergodic measure preserving transformations*, J. Eur. Math. Soc. (JEMS) 6 (2004), no. 3, 277-292.

[FM77] J. Feldman and C.C. Moore, *Ergodic equivalence relations and von Neumann algebras, I.*, Trans. Amer. Math. Soc, 234, 289-324, 1977.

[Ga00] D. Gaboriau, *Coût des relations d'équivalence et des groupes*, Invent. Math. 139 (2000), no. 1, 41-98.

[Ga02] D. Gaboriau, *Invariants* l^2 *de relations d'équivalence et de groupes.* Publ. Math. Inst. Hautes Études Sci. No. 95 (2002), 93-150.

[Gl12] Y. Glasner *Invariant random subgroups of Linear Groups*, preprint.

[HP11] P. de la Harpe, C^**-simple groups: amalgamated free products, HNN extensions, and fundamental groups of 3-Manifolds*, J. Topol. Anal. 3 (2011), no. 4, 451-489.

[Ha07] P. de la Harpe, *On simplicity of reduced* C^**-algebras of groups*, Bull. Lond. Math. Soc. 39 (2007), no. 1, 1-26.

[Ha85] P. de la Harpe, *Reduced* C^**-algebras of discrete groups which are simple with a unique trace.* Lecture Notes in Mathematics, 1132 (Springer-Verlag, 1985).

[Hu66] A. Hulanicki, *Means and Følner condition on locally compact groups*, Studia Math. 27, 1966, 87-104.

[Hu64] A. Hulanicki, *Groups whose regular representation weakly contains all unitary representations*, Studia Math. 24 1964 37-59.

[Io10] A. Ioana, *Non-orbit equivalent actions of* $\mathbb{F}_n$, Ann. Sci. Éc. Norm. Supér. (4) 42 (2009), no. 4, 675-696.

[Io11] A. Ioana, *Orbit inequivalent actions for groups containing a copy of* $\mathbb{F}_2$, Invent. Math. 185 (2011), no. 1, 55-73.

[IKT09] A. Ioana, A. Kechris, and T. Tsankov, *Subequivalence relations and positive definite functions*, Groups Geom. Dyn. 3 (2009), no. 4, 579-625.

[JKL02] S. Jackson, A. Kechris, and A. Louveau, *Countable Borel equivalence relations.* J. Math. Log. 2 (2002), no. 1, 1-80.

[Ke10] A. Kechris, *Global aspects of ergodic group actions*, Mathematical Surveys and Monographs, 160. American Mathematical Society, Providence, RI, 2010.

[Ke07] A. Kechris, *Unitary representations and modular actions*, J. Math. Sci. (N. Y.) 140 (2007), no. 3, 398-425.

[Ke95] A. Kechris, *Classical descriptive set theory*, Graduate Texts in Mathematics, 156. Springer-Verlag, New York, 1995.

[KM05] A. Kechris, B. Miller, *Topics in orbit equivalence.* Lecture Notes in Mathematics, 1852. Springer-Verlag, Berlin, 2004.

[KT09] A. Kechris, and T. Tsankov, *Amenable actions and almost invariant sets*, Proc. Amer. Math. Soc. 136 (2008), no. 2, 687-697.

[MOY11] Y. Matsuda, S.I. Oguni, S. Yamagata, C^**-simplicity for groups with non-elementary convergence group actions*, arXiv:1106.2618v2, 2011.

[OW80] D. Ornstein, and B. Weiss, *Ergodic theory of amenable group actions. I. The Rohlin lemma*, Bull. Amer. Math. Soc. (N.S.) 2 (1980), no. 1, 161-164.

[PS79] W. Paschke, and N. Salinas, C^**-algebras associated with free products of groups.* Pacific J. Math. 82 (1979), no. 1, 211-221.

[PT11] J. Peterson, and A. Thom, *Group cocycles and the ring of affiliated operators*, Invent. Math. 185 (2011), no. 3, 561-592.

[Pop08] S. Popa, *On the superrigidity of malleable actions with spectral gap*, J. Amer. Math. Soc. 21 (2008), no. 4, 981-1000.

[Pow75] R. Powers, *Simplicity of the* C^**-algebra associated with the free group on two generators*, Duke Math. J. 42 (1975) 151-156

[Poz09] T. Poznansky, *Characterization of linear groups whose reduced* C^**-algebras are simple*, arXiv:0812.2486v7. 2009.

[Pré12] J.P. Préaux, *Group extensions with infinite conjugacy classes*, arXiv:1201.4107v1, 2012.

[Pro93] S. Promislow, *A class of groups producing simple, unique trace,* C**-algebras*, Math. Proc. Cambridge Philos. Soc. 114 (1993) 223-233.

[ST10] R. Sauer and A. Thom, *A spectral sequence to compute* L^2*-Betti numbers of groups and groupoids*, J. Lond. Math. Soc. (2) 81 (2010), no. 3, 747-773.

[Sc81] K. Schmidt, *Amenability, Kazhdan's property T, strong ergodicity and invariant means for ergodic group-actions.* Ergodic Theory Dynamical Systems 1 (1981), no. 2, 223-236

[T-D12a] R. Tucker-Drob, *Weak equivalence and non-classifiability of measure preserving actions*, arXiv:1202.3101v2, 2012.

[T-D12b] R. Tucker-Drob, *Mixing actions of countable groups are almost free*, arXiv:1208.0655v1, 2012.

[Ta50] M. Takahasi, *Note on locally free groups.* J. Inst. Polytech. Osaka City Univ. Ser. A. Math. 1, (1950). 65-70.

[Ve11] A. Vershik, *Nonfree actions of countable groups and their characters*, J. Math. Sci. (N. Y.) 174 (2011), no. 1, 1-6.

[Ve12] A. Vershik, *Totally nonfree actions and the infinite symmetric group*, Mosc. Math. J. 12 (2012), no. 1, 193-212.

[Zi84] R. J. Zimmer. *Ergodic theory and semisimple groups.* Monographs in Mathematics, 81. Birkhauser Verlag, Basel, 1984.

Department of Mathematics
California Institute of Technology
Pasadena, CA 91125
`rtuckerd@caltech.edu`